

Research Article

Open Access

Dietmar Pfeifer*, Andreas Mändle, and Olena Ragulina

New copulas based on general partitions-of-unity and their applications to risk management (part II)

<https://doi.org/10.1515/demo-2017-0014>

Received March 16, 2017; accepted October 11, 2017

Abstract: We present a constructive and self-contained approach to data driven infinite partition-of-unity copulas that were recently introduced in the literature. In particular, we consider negative binomial and Poisson copulas and present a solution to the problem of fitting such copulas to highly asymmetric data in arbitrary dimensions.

Keywords: copulas, partition-of-unity, tail dependence, asymmetry

MSC: 62H05, 62H12, 62H17, 62H20

1 Introduction

Infinite partition-of-unity copulas have been introduced recently in the paper by Pfeifer et al. [9]. The main emphasis there was, however, on a particular symmetric case called diagonal dominance for which tail dependence coefficients could be explicitly calculated. The general asymmetric case was not treated in full detail. Our particular interest here is to complete the general setup with a suggestion how a data driven approach can be used to fit such copulas to highly asymmetric data in arbitrary dimensions, a question that had remained open so far.

2 A formal framework for infinite partition-of-unity copulas

Assume that $\{\varphi_{ki}(u)\}_{i \in \mathbb{Z}^+}$ for $k = 1, \dots, d \in \mathbb{N}$ represent discrete distributions over \mathbb{Z}^+ with a parameter $u \in (0, 1)$, i.e.

$$\varphi_{ki}(u) \geq 0 \quad \text{and} \quad \sum_{i=0}^{\infty} \varphi_{ki}(u) = 1 \quad \text{for } u \in (0, 1),$$

with

$$\alpha_{ki} := \int_0^1 \varphi_{ki}(u) du > 0 \quad \text{for } i \in \mathbb{Z}^+.$$

Let further $\{p_{\mathbf{i}}\}_{\mathbf{i} \in \mathbb{Z}^{+d}}$ represent the distribution of an arbitrary discrete d -dimensional random vector \mathbf{Z} over \mathbb{Z}^{+d} where, for simplicity, we write $\mathbf{i} = (i_1, \dots, i_d)$, and

$$P(\mathbf{Z} = \mathbf{i}) = p_{\mathbf{i}}, \quad \mathbf{i} \in \mathbb{Z}^{+d}.$$

***Corresponding Author: Dietmar Pfeifer:** Carl von Ossietzky Universität Oldenburg, Germany, E-mail: dietmar.pfeifer@uni-oldenburg.de

Andreas Mändle: Carl von Ossietzky Universität Oldenburg, Germany, E-mail: andreas.maendle@uni-oldenburg.de

Olena Ragulina: Taras Shevchenko National University of Kyiv, Ukraine, E-mail: ragulina.olena@gmail.com

Suppose further that for the marginal distributions, there holds

$$P(Z_k = i) = \alpha_{ki}, \quad i \in \mathbb{Z}^+, \quad k = 1, \dots, d.$$

Then

$$c(\mathbf{u}) := \sum_{\mathbf{i} \in \mathbb{Z}^{+d}} \frac{p_{\mathbf{i}}}{\prod_{k=1}^d \alpha_{k,i_k}} \prod_{k=1}^d \varphi_{k,i_k}(u_k), \quad \mathbf{u} = (u_1, \dots, u_d) \in (0, 1)^d \tag{1}$$

defines the density of a d -variate copula, which is called *infinite partition-of-unity copula* (IPU-copula for short).

Alternatively, we can rewrite (1) as

$$c(\mathbf{u}) = \sum_{\mathbf{i} \in \mathbb{Z}^{+d}} p_{\mathbf{i}} \prod_{k=1}^d f_{k,i_k}(u_k), \quad \mathbf{u} = (u_1, \dots, u_d) \in (0, 1)^d \tag{2}$$

where the $f_{ki}(\bullet) = \frac{\varphi_{ki}(\bullet)}{\alpha_{ki}}, i \in \mathbb{Z}^+, k = 1, \dots, d$ denote the Lebesgue densities induced by the $\{\varphi_{ki}(u)\}_{i \in \mathbb{Z}^+}$. A stochastic representation of the probability distribution induced by (1) or (2) is given by the random vector $\mathbf{U}(\mathbf{Z}) = \mathbf{U}(Z_1, \dots, Z_d) := U_{1,Z_1}, \dots, U_{d,Z_d}$ (with \mathbf{Z} as above) with stochastically independent random variables $\{U_{ki} | k = 1, \dots, d, i \in \mathbb{Z}^+\}$ (also independent of \mathbf{Z}) where the distribution of U_{ki} is induced by the density $f_{ki}(\bullet) = \frac{\varphi_{ki}(\bullet)}{\alpha_{ki}}, i \in \mathbb{Z}^+, k = 1, \dots, d$. \mathbf{Z} is called the *driver* of the IPU copula with density given in (1).

The following two classes of IPU copulas have been investigated in detail in [9], among others:

Example 1 (negative binomial copula). Let, for fixed integers $a_k > 0$,

$$\varphi_{ki}^{NB}(u) = \binom{a_k + i - 1}{i} u^i (1 - u)^{a_k} \text{ for } k, i \in \mathbb{Z}^+ \text{ and } u \in (0, 1). \tag{3}$$

Here we have $\alpha_{ki}^{NB} = \int_0^1 \varphi_{ki}^{NB}(u) du = \frac{a_k}{(a_k + i)(a_k + i + 1)}$ which corresponds to a discrete analogy of a Pareto distribution. The densities f_{ki}^{NB} are those of a beta distribution with parameters $(i+1, a_k+1)$. The corresponding copula density is thus given by

$$c^{NB}(\mathbf{u}) = \prod_{k=1}^d (a_k + 1) \sum_{\mathbf{i} \in \mathbb{Z}^{+d}} p_{\mathbf{i}} \prod_{k=1}^d \binom{a_k + i_k + 1}{i_k} u_k^{i_k} (1 - u_k)^{a_k}, \quad \mathbf{u} \in (0, 1)^d \tag{4}$$

with $p_{i_k} = P(Z_k = i_k) = \frac{a_k}{(a_k + i_k)(a_k + i_k + 1)}, i_k \in \mathbb{Z}^+, k = 1, \dots, d$.

Example 2 (Poisson copula). Let, for $L(u) := -\ln(1 - u) > 0, u \in (0, 1)$ and fixed parameters $a_k > 0$,

$$\varphi_{ki}^P(u) = (1 - u)^{a_k} \frac{a_k^i L(u)^i}{i!}. \tag{5}$$

Here we have $\alpha_{ki}^P = \int_0^1 \varphi_{ki}^P(u) du = \left(\frac{a_k}{a_k + 1}\right)^i \left(1 - \frac{a_k}{a_k + 1}\right)$, representing geometric distributions over \mathbb{Z}^+ . The densities f_{ki}^P are those of a transformed gamma distribution with $f_{ki}(u) = (a_k + 1)^{i+1} \frac{L^i(u)}{i!} (1 - u)^{a_k}$ for $k, i \in \mathbb{Z}^+$ and $u \in (0, 1)$. The corresponding copula density is thus given by

$$c^P(\mathbf{u}) = \prod_{k=1}^d (a_k + 1) \sum_{\mathbf{i} \in \mathbb{Z}^{+d}} p_{\mathbf{i}} \prod_{k=1}^d \frac{(a_k + 1)^k}{i_k!} L^{i_k}(u_k) (1 - u_k)^{i_k}, \quad \mathbf{u} \in (0, 1)^d \tag{6}$$

with $p_{i_k} = P(Z_k = i_k) = \left(\frac{a_k}{a_k + 1}\right)^{i_k} \left(1 - \frac{a_k}{a_k + 1}\right) = \frac{a_k^{i_k}}{(a_k + 1)^{i_k + 1}}, i_k \in \mathbb{Z}^+, k = 1, \dots, d$.

Note that a random variable X_{ki} with density f_{ki}^P can be represented as $X_{ki} = 1 - \exp(-Y_{ki})$, where Y_{ki} follows a gamma distribution with shape parameter $i + 1$ and scale parameter $a_k + 1$.

In [9], essentially the symmetrical case, i.e. the case of identical components of the driver \mathbf{Z} was considered (so called *diagonal dominance*). This means that the copula induced by the d -variate distribution of \mathbf{Z} is given by the upper Fréchet bound $C(\mathbf{u}) = \min(u_1, \dots, u_d)$, $\mathbf{u} \in (0, 1)^d$. In the two-dimensional case, the above formulas simplify to a great extent. In particular, it was proved that the negative binomial copula for arbitrary $a_1 = a_2 = a > 0$ has a positive tail dependence coefficient given by

$$\lambda_U(a) = \lim_{t \uparrow 1} \frac{\int_t^1 \int_t^1 c^{NB}(u, v) du dv}{1-t} = 1 - \frac{\binom{2a}{a}}{4^a} \sim 1 - \frac{1}{\sqrt{\pi a}} \text{ for large } a. \tag{7}$$

The Poisson copula, in contrast, has no tail dependence for all choices $a_1 = a_2 = a > 0$ although the density given by (6) is unbounded and has a pole in the point $(u, v) = (1, 1)$ (see Figures 1–2).




Figure 1: Density of a bivariate negative binomial copula, $\alpha = 5$




Figure 2: Density of a bivariate Poisson copula, $\alpha = 5$

So far the question remained open how such copulas could be fitted to highly *asymmetric* data in arbitrary dimensions. In the sequel, we shall give a constructive answer to this problem.

3 Constructing infinite partition-of-unity copulas from given data

The general idea here is to relate the $\{p_i\}_{i \in \mathbb{Z}^+}$, which essentially determine the structure of the IPU copula, to the empirical copula given by the data observed in an appropriate way. Let, for this purpose, denote \hat{C} a copula that is suitably estimated from the empirical copula, like a Bernstein copula, a rook copula (cf. [1]), appropriate shuffles of M (cf. [7, Section 3.2.3]) or other patchwork copulas that can be easily simulated by Monte Carlo methods (cf. [2] or [3]). Let further denote F_k the cumulative distribution function induced by the discrete distribution $\alpha_{k\bullet}$, i.e. $F_k(i) = \sum_{j=0}^i \alpha_{kj}$, $i \in \mathbb{Z}^+$. If $\hat{\mathbf{U}} = (\hat{U}_1, \dots, \hat{U}_d)$ denotes a stochastic representation of \hat{C} , set $Z_k := F_k^{-1}(\hat{U}_k)$ for $k = 1, \dots, d$. Then $\mathbf{Z} = (Z_1, \dots, Z_d)$ is an appropriate driver for a data-driven IPU copula.

For the examples given above, the resulting formulas are quite simple.

Example 1: $Z_k = F_k^{-1}(\hat{U}_k) = \lfloor \frac{\alpha_k \hat{U}_k}{1 - \hat{U}_k} \rfloor$, where $\lfloor z \rfloor = \max\{x \in \mathbb{R} \mid x \leq z\}$ (round down).

Example 2: $Z_k = F_k^{-1} \hat{U}_k = \left\lfloor \frac{-\ln(1 - \hat{U}_k)}{\ln(a_k + 1) - \ln a_k} \right\rfloor$.

This follows immediately from standard arguments in Monte Carlo theory: in Example 1, we have

$$Z_k = \left\lfloor \frac{a_k \hat{U}_k}{1 - \hat{U}_k} \right\rfloor = i \quad \text{iff} \quad \frac{i}{a_k + i} \leq \hat{U}_k < \frac{i + 1}{a_k + i + 1},$$

with probabilities

$$P(Z_k = i) = P\left(\frac{i}{a_k + i} \leq \hat{U}_k < \frac{i + 1}{a_k + i + 1}\right) = \frac{a_k}{(a_k + i)(a_k + i + 1)}$$

for $i \in \mathbb{Z}^+$, as desired.

In Example 2, we have

$$Z_k = F_k^{-1} \hat{U}_k = \left\lfloor \frac{-\ln(1 - \hat{U}_k)}{\ln(a_k + 1) - \ln a_k} \right\rfloor = i \quad \text{iff} \quad 1 - \left(\frac{a_k}{a_k + 1}\right)^i \leq \hat{U}_k < 1 - \left(\frac{a_k}{a_k + 1}\right)^{i+1},$$

with probabilities

$$P(Z_k = i) = P\left(1 - \left(\frac{a_k}{a_k + 1}\right)^i \leq \hat{U}_k < 1 - \left(\frac{a_k}{a_k + 1}\right)^{i+1}\right) = \frac{a_k^i}{(a_k + 1)^{i+1}} = \left(\frac{a_k}{a_k + 1}\right)^i \left(1 - \frac{a_k}{a_k + 1}\right)$$

for $i \in \mathbb{Z}^+$, as desired.

The method proposed here allows for a great flexibility concerning the construction of data-driven IPU copulas, including cases with positive tail dependence. We discuss this here along the example given in [1], Example 4.2, which was also the basis for the discussion in [9, Section 4].

Table 1 and Figures 3–4 show the original data (x_i, y_i) and the corresponding rank vectors (r_{1i}, r_{2i}) .

Table 1: The original data and the corresponding rank vectors

i	x_i	y_i	r_{1i}	r_{2i}
1	0.468	0.966	4	9
2	9.951	2.679	20	20
3	0.866	0.897	8	4
4	6.731	2.249	19	19
5	1.421	0.956	13	8
6	2.040	1.141	17	15
7	2.967	1.707	18	18
8	1.200	1.008	11	10
9	0.426	1.065	3	12
10	1.946	1.162	15	16
11	0.676	0.918	5	6
12	1.184	1.336	10	17
13	0.960	0.933	9	7
14	1.972	1.077	16	13
15	1.549	1.041	14	11
16	0.819	0.899	6	5
17	0.063	0.710	1	1
18	1.280	1.118	12	14
19	0.824	0.894	7	3
20	0.227	0.837	2	2

The graphs in Figures 5–14 show 5,000 Monte Carlo simulations each from different constructions of data-driven IPU copulas (small dots), with a superposition of the empirical copula (scaled rank vectors) as large




Figure 3: Graph of original data




Figure 4: Graph of rank vectors

white points. The symmetric cases (negative binomial and Poisson copulas) were constructed according to the suggestions in [9, Section 4]. The asymmetric cases were constructed on the basis of a shuffle of M copula with local upper Fréchet bounds for the driver Z , shown first. Note that the location of the corresponding line sections are one to one determined by the relative rank vectors from the original data.




Figure 5: Relative ranks $r_{.i}/20$




Figure 6: Shuffle of M with local upper Fréchet bounds

As can be clearly seen, the asymmetric IPU copula follows the given data (empirical copula) much better than the symmetric IPU copulas. Also, in contrast to the Bernstein copula with no tail dependence, the asymmetric negative binomial IPU copulas always show a positive tail dependence. Since tail dependence is an asymptotic property, we can conclude that the corresponding tail dependence coefficient in the above constructions can be calculated from the parameter $a = b$ according to relation (7). Note also that although the pictures above might suggest a positive tail dependence for the asymmetric Poisson IPU copula, this is theoretically not possible.

4 Implications for risk management

The new European supervisory regulations in the financial sector (Basel III for banks, Solvency II for insurance companies) require the calculation of a sufficient capital adequacy based on the risk measure




Figure 7: Bernstein copula




Figure 8: Symmetric negative binomial copula, $a = b = 5$




Figure 9: Asymmetric negative binomial copula, $a = b = 5$




Figure 10: Asymmetric negative binomial copula, $a = b = 10$




Figure 11: Symmetric Poisson copula, $a = b = 6$




Figure 12: Asymmetric Poisson copula, $a = b = 6$




Figure 13: Asymmetric Poisson copula,
 $a = b = 10$




Figure 14: Asymmetric Poisson copula,
 $a = b = 15$

Value@Risk $\text{VaR}_\alpha(S)$, which is defined as the $1 - \alpha$ quantile of the distribution of the total portfolio risk $S = \sum_{i=1}^d X_i$ where X_1, \dots, X_d are the individual risk positions in the portfolio. Especially in internal models for the calculation of the underlying aggregate risk measure, it is important to find appropriate models for the stochastic dependence between individual risk positions. It is well known that in the case of comonotonicity between risks – i.e., the underlying copula is the upper Fréchet bound – there is no diversification effect and the risk measure Value@Risk is additive (cf. [6, Proposition 7.20]). Also, the worst case for the total Value@Risk is not attained under comonotonicity but rather in cases where there is a kind of local negative dependence in the upper right corner of the underlying copula (cf. [10] or [4]). A similar negative result holds for an assumed dependence between correlation and diversification (cf. [8]), which is frequently stated in the common legislative papers. The following examples show how the asymmetric data-driven IPU copula approach can provide competing estimates for the risk measure Value@Risk on the basis of the same data observed. For the sake of simplicity, we use the data set from [1], Example 4.2, discussed above. We compare the following IPU copula approaches:

- a classical Bernstein copula with grid size 20 (cf. [1])
- the asymmetric negative binomial copula with parameters $a = b = 5$, $a = b = 10$ and $a = b = 15$ (for short: NB 5, NB 10 and NB 15)
- the asymmetric Poisson copula with parameters $a = b = 6$, $a = b = 10$ and $a = b = 15$ (for short: Po 6, Po 10 and Po 15)
- "worst case" (WC) versions of these copulas where a particular shuffle of M is used (i.e. with a local lower Fréchet bound in the upper right corner)

The graphs in Figures 15–16 show the corresponding "worst case" copula driver. The graphs in Figures 17–22 show scatterplots from 5,000 simulations each for the underlying "worst case" copulas.

The estimates in Tables 2–3 are based on 5,000,000 Monte Carlo simulations for each particular copula approach. For the marginal distributions of the two risk positions X and Y , a lognormal and a Fréchet distribution were estimated from the original data. The risk level α was chosen as $\alpha = 0.05$. With the estimated parameters for the marginal distributions, we have $\widehat{\text{VaR}}_{0.05}(X) = 6.8190$, $\widehat{\text{VaR}}_{0.05}(Y) = 2.0984$ and $\widehat{\text{VaR}}_{0.05}(X) + \widehat{\text{VaR}}_{0.05}(Y) = 8.9174$. From the simulations, we obtain, with $S = X + Y$, results presented in Tables 2–3.

The graphs in Figure 23 show some empirical quantile functions of the simulations above.

These results clearly show that the "worst case" IPU copula approaches always result in a risk concentration effect while the basic negative binomial and the Poisson copula approach show a slight diversification effect which decreases with increasing parameters $a = b$. Note that the results for the negative binomial and the Poisson copula are quite close in spite of the fact that the negative binomial copula here always has a




Figure 15: Relative ranks $r_i/20$




Figure 16: "Worst case" shuffle of M




Figure 17: "Worst case" negative binomial copula, $a = b = 5$




Figure 18: "Worst case" negative binomial copula, $a = b = 10$




Figure 19: "Worst case" negative binomial copula, $a = b = 15$




Figure 20: "Worst case" Poisson copula, $a = b = 6$




Figure 21: "Worst case" Poisson copula, $a = b = 10$




Figure 22: "Worst case" Poisson copula, $a = b = 15$

Table 2: Estimates based on Monte Carlo simulations. Bernstein and negative binomial copula approaches

Copula type	Bernstein	NB 5	NB 5 WC	NB 10	NB 10 WC	NB 15	NB 15 WC
$\widehat{\text{VaR}}_{0.05}(S)$	8.9586	8.8474	9.3989	8.8834	9.5421	8.8978	9.6198
$\widehat{\text{VaR}}_{0.05}(X) + \widehat{\text{VaR}}_{0.05}(Y)$	8.9174	8.9174	8.9174	8.9174	8.9174	8.9174	8.9174

Table 3: Estimates based on Monte Carlo simulations. Poisson copula approaches

Copula type	Po 6	Po 6 WC	Po 10	Po 10 WC	Po 15	Po 15 WC
$\widehat{\text{VaR}}_{0.05}(S)$	8.8200	9.1402	8.8453	9.2412	8.8820	9.3532
$\widehat{\text{VaR}}_{0.05}(X) + \widehat{\text{VaR}}_{0.05}(Y)$	8.9174	8.9174	8.9174	8.9174	8.9174	8.9174





Figure 23: Empirical quantile functions

positive tail dependence. It is interesting to see that the Bernstein copula also shows a risk concentration effect although there is no tail dependence and also no strict "worst case" behaviour.

5 Conclusion

The IPU copula approach for asymmetric data sets is a very flexible tool to model dependencies between risks, also in higher dimensions. It covers cases of tail dependence and also of "worst case" scenarios on the basis of the same data set. It follows the shape of the data more closely than most other approaches and can easily be implemented in usual spreadsheets. Note that our motivation for a patchwork construction for the copula driver resembles very much the arguments in [3]. The difference is, however, that the resulting IPU copula itself is not a patchwork copula.

Especially in the light of the new European supervisory regulations in the financial sector such approaches might be interesting to figure out unfavourable constellations which lead to a higher demand of equity or solvency capital. It should be kept in mind, however, that this is not only a problem of the assumed underlying copula, but also depends significantly on the type of the marginal risk distributions, as is discussed in [5].

Acknowledgement: We would like to thank the referees for some critical comments on an earlier version which lead to a clearer exposition of the content of the paper.

References

- [1] Cottin, C. and D. Pfeifer (2014). From Bernstein polynomials to Bernstein copulas. *J. Appl. Funct. Anal.* 9(3-4), 277–288.
- [2] Durante, F. and J. Fernández-Sánchez (2010). Multivariate shuffles and approximation of copulas. *Statist. Probab. Lett.* 80(23-24), 1827–1834.
- [3] Durante, F., J. Fernández-Sánchez, and C. Sempi (2013). Multivariate patchwork copulas: a unified approach with applications to partial comonotonicity. *Insur. Math. Econ.* 53(3), 897–905.
- [4] Embrechts, P., G. Puccetti, and L. Rüschendorf (2013). Model uncertainty and VaR aggregation. *J. Bank. Financ.* 37(8), 2750–2764.
- [5] Ibragimov, R. and A. Prokhorov (2017). *Heavy Tails and Copulas: Topics in Dependence Modelling in Economics and Finance*. World Scientific, Singapore.
- [6] McNeil, A. J., R. Frey, and P. Embrechts (2015). *Quantitative Risk Management: Concepts, Techniques and Tools*. Revised edition. Princeton University Press.
- [7] Nelsen, R. B. (2006). *An Introduction to Copulas*. Second edition. Springer, New York.
- [8] Pfeifer, D. (2013). Correlation, tail dependence and diversification. In C. Becker, R. Fried, and S. Kuhnt (Eds.), *Robustness and Complex Data Structures: Festschrift in Honour of Ursula Gather*, pp. 301–314. Springer, Berlin.
- [9] Pfeifer, D., H. A. Tsatedem, A. Mändle, and C. Girschig (2016). New copulas based on general partitions-of-unity and their applications to risk management. *Depend. Model.* 4, 123–140.
- [10] Puccetti, G. and L. Rüschendorf (2012). Computation of sharp bounds on the distribution of a function of dependent risks. *J. Comput. Appl. Math.* 236(7), 1833–1840.