

On Neumann-Poincaré operators and self-adjoint transmission problems

Badreddine Benhellal † and Konstantin Pankrashkin ‡

Carl von Ossietzky Universität Oldenburg
Fakultät V – Mathematik und Naturwissenschaften
Institut für Mathematik
26111 Oldenburg, Germany

† E-Mail: badreddine.benhellal@uol.de

‡ ORCID: 0000-0003-1700-7295

E-Mail: konstantin.pankrashkin@uol.de,

Webpage: <http://uol.de/pankrashkin>

Abstract

We discuss the self-adjointness in L^2 -setting of the operators acting as $-\nabla \cdot h \nabla$, with piecewise constant functions h having a jump along a Lipschitz hypersurface Σ , without explicit assumptions on the sign of h . We establish a number of sufficient conditions for the self-adjointness of the operator with $H^{\frac{3}{2}}$ -regularity in terms of the jump value and the regularity and geometric properties of Σ . An important intermediate step is a link with Fredholm properties of the Neumann-Poincaré operator on Σ , which is new for the Lipschitz setting.

1 Introduction

Let $n \geq 2$ and $\Omega \subset \mathbb{R}^n$ be a non-empty bounded open set with Lipschitz boundary. Let $\Omega_- \subset \Omega$ be a non-empty open subset with Lipschitz boundary such that $\overline{\Omega_-} \subset \Omega$, and set

$$\Omega_+ = \Omega \setminus \overline{\Omega_-}, \quad \Sigma = \partial\Omega_-, \quad (1)$$

see Fig. 1. In order to avoid combinatorially involved configurations we assume from the very beginning that

Σ is connected.




Figure 1: The sets Ω_{\pm} and Σ .

In the present work, we will be interested in the self-adjointness of the operator formally acting as $u \mapsto -\nabla \cdot h \nabla u$ in $L^2(\Omega)$ with the Dirichlet boundary conditions at the exterior boundary $\partial\Omega$, where $h : \Omega \rightarrow \mathbb{R}$ is the piecewise constant function

$$h : \Omega \ni x \mapsto \begin{cases} 1, & x \in \Omega_+, \\ \mu, & x \in \Omega_- \end{cases}$$

and $\mu \in \mathbb{R} \setminus \{0\}$ is a parameter. For $\mu > 0$ the operator can be defined using Lax-Milgram representation theorem and the ellipticity theory, and it is semibounded from below with compact resolvent, so we are mainly interested in the case $\mu < 0$. Various equations with the above operators for $\mu < 0$ appear in the mathematical theory of negative-index metamaterials, stemming from the pioneering works by Veselago [43], and they show a number of unexpected properties compared to the usual elliptic case [13, 26]. We also remark that there is an increasing interest to sign-changing problems in the homogenization theory and the uncertainty quantification [33, 34, 44].

In what follows, any function $u \in L^2(\Omega)$ will be identified with the pair (u_+, u_-) with u_{\pm} being its restrictions on Ω_{\pm} . This gives rise to the identification $L^2(\Omega) \simeq L^2(\Omega_+) \oplus L^2(\Omega_-)$. Then the above operator is expected to act, at least at a naive level, as $(u_+, u_-) \mapsto (-\Delta u_+, -\mu \Delta u_-)$ while the functions u_{\pm} should satisfy the boundary and transmission conditions

$$u_+ = 0 \text{ on } \partial\Omega, \quad u_+ = u_- \text{ on } \Sigma, \quad \partial_+ u_+ + \mu \partial_- u_- \text{ on } \Sigma, \quad (2)$$

with ∂_{\pm} being the outward normal (with respect to Ω_{\pm}) derivative on Σ . Finding precise regularity properties of the functions u_{\pm} guaranteeing the self-adjointness of the resulting operator is a non-trivial task depending on a combination of the parameter μ (usually called contrast) and the geometric properties of Σ , and this is the central topic of the present paper.

It seems that the problem was first addressed in [12] for the case when both Ω_{\pm} are C^2 -smooth. Using the results of [16] it turned out that the

requirement $u_{\pm} \in H^1(\Omega_{\pm})$ with $\Delta u_{\pm} \in L^2(\Omega_{\pm})$ leads to a self-adjoint operator for all $\mu \neq -1$. On the other hand, if $n = 2$ and Σ has corners, then the operator is only self-adjoint if μ lies outside some non-trivial critical interval, see Remark 21 below. A similar picture was found for some boundary singularities in higher-dimensional situations as well, see [9–11].

It should be noted that the regularity of functions in the operator domain turns out to be crucially important. For example, in the papers [4, 14, 38] it was found that for some values of μ the operator with $u_{\pm} \in H^2(\Omega_{\pm})$ is not closed; its closure is a self-adjoint operator whose essential spectrum can be non-empty, which in parts explains various regularization issues found in earlier works [13, 25, 26]. We remark that very similar effects were recently discovered in transmission problems for Dirac operators as well [3, 7, 8].

It seems that no previous work addressed the case of general Lipschitz Σ so far, and we are trying to close this gap in the present paper. One should mention the existence of an approach based on modified representation theorems for quadratic forms [22, 27, 39, 40], which allows for the study of variable coefficients, but the final results on the self-adjointness require higher regularity of the hypersurface Σ or suitable symmetries of the sets Ω_{\pm} .

More precisely, denote by A the linear operator in $L^2(\Omega)$ acting as $(u_+, u_-) \mapsto (-\Delta u_+, -\mu \Delta u_-)$ on the functions $u_{\pm} \in H^{\frac{3}{2}}(\Omega_{\pm})$ such that $\Delta u_{\pm} \in L^2(\Omega_{\pm})$ and that the boundary and transmission conditions (2) are satisfied (a precise definition of the boundary values will be given in Sections 2 and 3). Our main result is a link between the self-adjointness of A and the spectral properties of the so-called Neumann-Poincaré operator K_{Σ} on Σ . Recall that K_{Σ} is a bounded linear operator $K_{\Sigma} : L^2(\Sigma) \rightarrow L^2(\Sigma)$ defined by

$$K_{\Sigma}f(x) = \text{p.v.} \int_{\Sigma} \frac{\langle \nu(y), y - x \rangle}{\sigma_n |x - y|^n} f(y) \, ds(y),$$

where ν is the unit normal field of Σ pointing to Ω_+ and p.v. stands for the Cauchy principal value. (We remark that the terminology is not unique: in some references, it is the adjoint of K_{Σ} which is referred to as the Neumann-Poincaré operator). The Neumann-Poincaré operator appears naturally when considering the boundary values of the double layer potential, see Section 2. Its spectral analysis attracted a lot of attention during the last decades [1, 15, 23, 28, 31, 32], and it already appeared in various problems related to the existence of solutions in sign-indefinite problems [21, 35–37]. Nevertheless, it seems that the links between the self-adjointness of A and the spectrum of K_{Σ} were never addressed explicitly for the case of general

Lipschitz Σ . Our main observation is as follows (Corollary 16):

if $\mu \in \mathbb{R} \setminus \{-1, 0\}$ and $K_\Sigma - \frac{\mu+1}{2(\mu-1)}$ is Fredholm of index zero, then A is self-adjoint.

Using this main condition, we establish several specific situations in which A is self-adjoint:

- the almost trivial case $\mu > 0$ (Theorem 17),
- for all $\mu \neq -1$, if ν has a vanishing mean oscillation, which covers all C^1 -smooth Σ (Theorem 18),
- $n = 2$, Σ is a curvilinear polygon with C^1 -smooth edges and the sharpest corner $\omega \in (0, \pi)$, and

$$\mu \notin \left[-\cot^2 \frac{\omega}{4}, -\tan^2 \frac{\omega}{4} \right].$$

The compatibility of the last condition with the earlier conditions from [12, 17] is discussed in Remark 21.

As discussed in Remark 19, the required spectral bounds for K_Σ belong actually to the most important conjectures in the theory of Neumann-Poincaré operators. Very few explicit results are available so far, which in part explains why the self-adjointness problem for the sign-indefinite case turns out to be unexpectedly difficult. We hope that the link between the two classes of problems we establish in this paper will draw the attention of new communities to the respective questions.

Our approach follows the philosophy of quasi boundary triples for symmetric operators [5, 6], which is enhanced by the very recent sharp trace theory on Lipschitz domains developed in [2]. Our main step is establishing a resolvent formula for the operator A (Proposition 13). Then the self-adjointness of A is reduced to proving the surjectivity of $A - z$ for non-real z , which is established with the help of the mapping properties of various boundary integral operators [42].

2 Boundary integral operators and Dirichlet-to-Neumann maps in Lipschitz domains

Let $n \geq 2$ and $U \subset \mathbb{R}^n$ be a non-empty open set with Lipschitz boundary ∂U and ν be the outer unit normal on ∂U .

We denote by H^s the usual Sobolev spaces of order $s \in \mathbb{R}$. In addition, we denote

$$H_\Delta^s(U) := \{f \in H^s(U) : \Delta f \in L^2(U)\},$$

which will be equipped with the norm $\|f\|_{H^s_\Delta(U)}^2 := \|f\|_{H^s(U)}^2 + \|\Delta f\|_{L^2(U)}^2$.

Let $\gamma_D^{\partial U} : H^1(U) \rightarrow L^2(\partial U)$ be the Dirichlet trace map, i.e.

$$\gamma_D^{\partial U} f := f|_{\partial U} \text{ for } f \in C^\infty(\bar{U}),$$

which is extended by density and continuity. In fact, better mapping properties are known [2, Corollary 3.7]:

Proposition 1. *The map $\gamma_D^{\partial U} : H^{\frac{3}{2}}_\Delta(U) \rightarrow H^1(\partial U)$ is bounded and surjective. There exists a bounded linear map $\Upsilon : H^1(\partial U) \rightarrow H^{\frac{3}{2}}_\Delta(U)$ such that*

$$\Delta \Upsilon \varphi = 0 \text{ and } \gamma_D^{\partial U} \Upsilon \varphi = \varphi \text{ for any } \varphi \in H^1(\partial U).$$

In addition, let $\gamma_N^{\partial U} : H^{\frac{3}{2}}_\Delta(U) \rightarrow L^2(\partial U)$ be the Neumann trace,

$$\gamma_N^{\partial U} f := \langle \nu, \nabla f|_{\partial U} \rangle \text{ for } f \in C^\infty(\bar{U}),$$

extended by density and continuity. The following properties are known [2, Corollary 5.7]:

Proposition 2. *The above map $\gamma_N^{\partial U}$ is well-defined, bounded and surjective, and for any $f, g \in H^{\frac{3}{2}}_\Delta(U)$ one has the usual Green formula (integration by parts)*

$$\langle f, \Delta g \rangle_{L^2(U)} - \langle \Delta f, g \rangle_{L^2(U)} = \langle \gamma_D^{\partial U} f, \gamma_N^{\partial U} g \rangle_{L^2(\partial U)} - \langle \gamma_N^{\partial U} f, \gamma_D^{\partial U} g \rangle_{L^2(\partial U)}.$$

Denote by Φ the fundamental solution of the Laplace equation in \mathbb{R}^n defined by

$$\Phi : \mathbb{R}^n \setminus \{0\} \ni x \mapsto \begin{cases} \frac{1}{2\pi} \log |x|, & \text{for } n = 2, \\ \frac{1}{\sigma_n(n-2)|x|^{n-2}}, & \text{for } n \geq 3, \end{cases},$$

where σ_n stands for the surface measure of the unit sphere in \mathbb{R}^n . Define the single layer boundary operator $S_{\partial U} : L^2(\partial U) \rightarrow L^2(\partial U)$ by

$$S_{\partial U} f(\cdot) := \int_{\partial U} \Phi(\cdot - y) f(y) \, ds(y).$$

It is clear from the definition that $S_{\partial U} : L^2(\partial U) \rightarrow L^2(\partial U)$ is bounded and self-adjoint. In addition, $\text{ran } S_{\partial U} \subset H^1(\partial U)$ and

$$S_{\partial U} : L^2(\partial U) \rightarrow H^1(\partial U)$$

is a bounded operator as well, see [42, Lemma 1.8]. The following finer properties of $S_{\partial U}$ are also known:

Proposition 3. *Assume that ∂U is connected.*

(a) *If $n \geq 3$, then $S_{\partial U} : L^2(\partial U) \rightarrow H^1(\partial U)$ is bijective.*

(b) *If $n = 2$, then two cases are possible:*

(i) *$S_{\partial U}$ is injective. Then $S_{\partial U} : L^2(\partial U) \rightarrow H^1(\partial U)$ is bijective.*

(ii) *There exists $f_0 \in L^2(\partial U)$ such that*

$$\int_{\partial U} f_0 \, ds = 1, \quad \ker S_{\partial U} = \mathbb{C}f_0.$$

Denote

$$L_0^2(\partial U) := \left\{ f \in L^2(\partial U) : \int_{\partial U} f \overline{f_0} \, ds = 0 \right\},$$

then $S_{\partial U} : L_0^2(\partial U) \rightarrow H^1(\partial U) \cap L_0^2(\partial U)$ is injective.

The part (a) is shown in [42, Theorem 3.3]. The part (b.i) is shown in [42, Theorem 4.11]. The part (b.ii) is not stated explicitly in [42], but follows directly from the argument in the proof of [42, Theorem 4.11].

We define the double layer operator $K_{\partial U} : L^2(\partial U) \rightarrow L^2(\partial U)$,

$$K_{\partial U} f(\cdot) = \text{p.v.} \int_{\partial U} \frac{\langle \nu(y), y - \cdot \rangle}{|\sigma_n| \cdot |y - \cdot|^n} f(y) \, ds(y).$$

It is also known to be bounded, [42], and its adjoint $K_{\partial U}^*$ is given by

$$K_{\partial U}^* f(\cdot) = \text{p.v.} \int_{\partial U} \frac{\langle \nu(\cdot), \cdot - y \rangle}{|\sigma_n| \cdot |y - \cdot|^n} f(y) \, ds(y).$$

Definition 4. The Dirichlet Laplacian $-\Delta_U$ associated with U is the linear operator in $L^2(U)$ defined by

$$\text{dom}(-\Delta_U) := \{f \in H_{\Delta}^{\frac{3}{2}}(U) : \gamma_D^{\partial U} f = 0\}, \quad -\Delta_U : f \mapsto -\Delta f,$$

which is known to be a self-adjoint operator with compact resolvent [2, Theorem 6.9]. Remark that for any $z \in \mathbb{C} \setminus \sigma(-\Delta_U)$ the resolvent $(-\Delta_U - z)^{-1}$ defines a bounded operator $L^2(U) \rightarrow H_{\Delta}^{\frac{3}{2}}(U)$.

Lemma 5. *Let $\varphi \in H^1(\partial U)$ and $z \in \mathbb{C} \setminus \sigma(-\Delta_U)$, then the boundary value problem*

$$(-\Delta - z)f = 0 \text{ in } U, \quad \gamma_D^{\partial U} f = \varphi, \quad f \in H_{\Delta}^{\frac{3}{2}}(U),$$

has a unique solution, which is given by

$$f := z(-\Delta_U - z)^{-1} \Upsilon \varphi + \Upsilon \varphi$$

with Υ defined in Proposition 1. The resulting map $\varphi \mapsto f$, i.e.

$$\begin{aligned} P_z^U &:= z(-\Delta_U - z)^{-1}\Upsilon + \Upsilon \\ &\equiv \left[z(-\Delta_U - z)^{-1} + 1 \right] \Upsilon : H^1(\partial U) \rightarrow H_{\Delta}^{\frac{3}{2}}(U) \end{aligned}$$

will be called the Poisson operator for U and z . By construction, it is a bounded operator.

Proof. One can uniquely represent $f := g + \Upsilon\varphi$ with $g \in \text{dom}(-\Delta_U)$. The substitution into the problem shows that f is a solution if and only if $(-\Delta - z)g = -(-\Delta - z)\Upsilon\varphi$.

Due to the properties of Υ this is equivalent to $(-\Delta_U - z)g = z\Upsilon\varphi$. As $\Upsilon\varphi \in L^2(U)$ and $z \in \mathbb{C} \setminus \sigma(-\Delta_U)$, one has a unique solution g given by $g = z(-\Delta_U - z)^{-1}\Upsilon\varphi$. \square

Definition 6. For $z \in \mathbb{C} \setminus \sigma(-\Delta_U)$ we define the Dirichlet-to-Neumann map N_z^U by

$$N_z^U := \gamma_N^{\partial U} P_z^U : H^1(\partial U) \rightarrow L^2(\partial U).$$

By construction, it is again a bounded operator.

Proposition 7. For each $z \in \mathbb{C} \setminus \sigma(-\Delta_U)$ the operator

$$N_z^U - N_0^U : H^1(\partial U) \rightarrow L^2(\partial U)$$

is compact.

Proof. Using the above representation we have

$$N_z^U - N_0^U = \gamma_N^{\partial U} (P_z^U - P_0^U) = z\gamma_N^{\partial U} (-\Delta_U - z)^{-1}\Upsilon.$$

The embedding $J : H^{\frac{3}{2}}(U) \rightarrow L^2(U)$ is compact, and one can rewrite

$$N_z^U - N_0^U = z\gamma_N^{\partial U} (-\Delta_U - z)^{-1}J\Upsilon,$$

and the compactness follows from the boundedness of all the other factors. \square

The following relation will be important:

Proposition 8. There holds

$$S_{\partial U} N_0^U = \frac{1}{2} - K_{\partial U}. \quad (3)$$

Proof. For $g \in L^2(\partial U)$ and $x \in U$ define the single and double layer potentials,

$$\begin{aligned}\mathcal{S}g(x) &= \int_{\partial U} \Phi(x-y)g(y) \, ds(y), \\ \mathcal{D}g(x) &= \int_{\partial U} \frac{\langle \nu(y), y-x \rangle}{\sigma_n |x-y|^n} g(y) \, ds(y).\end{aligned}$$

It is known that $\mathcal{S} : L^2(\partial U) \rightarrow H^1(U)$ and $\mathcal{D} : L^2(\partial U) \rightarrow H^1(U)$ are well-defined and bounded, see [30, Corollary 5.2.9 and Theorem 5.6.1]. In addition, we have the following identities:

$$\gamma_D^{\partial U} \mathcal{S} = S_{\partial U}, \quad \gamma_N^{\partial U} \mathcal{S} = -\frac{1}{2} + K_{\partial U}^*, \quad \gamma_D^{\partial U} \mathcal{D} = \frac{1}{2} + K_{\partial U}, \quad (4)$$

see [29, Theorem 6.11 and Eq. (7.5)].

Let $\psi \in H^1(\partial U)$ and set $u = P_0^U \psi$. By [30, Theorem 5.6.1] we have

$$u = \mathcal{D}\gamma_D^{\partial U} u + \mathcal{S}\gamma_N^{\partial U} u \equiv \mathcal{D}\psi + \mathcal{S}N_0^U \psi.$$

Using (4) we get

$$\psi = \gamma_D^{\partial U} \mathcal{D}\psi + \gamma_D^{\partial U} \mathcal{S}N_0^U \psi = \left(\frac{1}{2} + K_{\partial U}\right)\psi + \mathcal{S}N_0^U \psi. \quad \square$$

Also, the following relation will be important:

$$S_{\partial U} K_{\partial U}^* = K_{\partial U} S_{\partial U}, \quad (5)$$

known as the symmetrization formula, [30, Proposition 5.7.1].

For subsequent use, we mention the following identity, which follows from the integration by parts:

Lemma 9. *For any $\varphi \in H^1(\partial U)$, any $z \in \mathbb{C} \setminus \sigma(-\Delta_U)$ and $f := P_z^U \varphi$ there holds*

$$\langle \varphi, N_z^U \varphi \rangle_{L^2(\partial U)} = \int_U |\nabla f|^2 \, dx - z \int_U |f|^2 \, dx.$$

In particular, N_z^U is a symmetric operator in $L^2(\partial U)$ for real z .

We discuss in greater detail the case when ∂U is disconnected. Let Λ be a connected component of ∂U and $\Lambda_0 := \partial U \setminus \Lambda \neq \emptyset$. This gives rise to decompositions

$$H^s(\partial U) \simeq H^s(\Lambda) \oplus H^s(\Lambda_0)$$

and similar decompositions for other functional spaces over U , as well as to the representation of trace maps $\gamma_{D/N}^{\partial U}$ as

$$\gamma_{D/N}^{\partial U} f = (\gamma_{D/N}^{\Lambda} f, \gamma_{D/N}^{\Lambda_0} f), \quad \gamma_{D/N}^{\Lambda} f := \gamma_{D/N}^{\partial U} f|_{\Lambda}, \quad \gamma_{D/N}^{\Lambda_0} f := \gamma_{D/N}^{\partial U} f|_{\Lambda_0}.$$

For $z \notin \sigma(-\Delta_U)$ we introduce the partial Dirichlet-to-Neumann operators

$$\tilde{N}_z^U : H^1(\Lambda) \ni \varphi \mapsto \gamma_N^\Lambda P_z^U(\varphi, 0) \in L^2(\Lambda),$$

i.e. $\tilde{N}_z^U \varphi := \gamma_N^\Lambda f$, where f is the unique solution to

$$(-\Delta - z)f = 0 \text{ in } U, \quad \gamma_D^\Lambda f = \varphi, \quad \gamma_D^{\Lambda_0} f = 0.$$

In view of the relation with N_z^U one obtains

Proposition 10. *For each $z \in \mathbb{C} \setminus \sigma(-\Delta_U)$ the operator $\tilde{N}_z^U - \tilde{N}_0^U : H^1(\Lambda) \rightarrow L^2(\Lambda)$ is compact.*

We will need the following variation of Proposition 8:

Proposition 11. *There holds*

$$S_\Lambda \tilde{N}_0^U = \frac{1}{2} - K_\Lambda + B,$$

where $B : H^1(\Lambda) \rightarrow L^2(\Lambda)$ is a compact operator.

Proof. The operators $K_{\partial U}$ and $S_{\partial U}$ are decomposed as

$$K_{\partial U} = \begin{pmatrix} K_\Lambda & K_{12} \\ K_{21} & K_{\Lambda_0} \end{pmatrix}, \quad S_{\partial U} = \begin{pmatrix} S_\Lambda & S_{12} \\ S_{21} & S_{\Lambda_0} \end{pmatrix}, \quad N_0^{\partial U} = \begin{pmatrix} \tilde{N}_0^U & N_{12} \\ N_{21} & N_{22} \end{pmatrix},$$

so taking the upper left corner in (3) gives

$$S_\Lambda \tilde{N}_0^U + S_{12} N_{21} = \frac{1}{2} - K_\Lambda,$$

where by construction S_{12} and N_{21} are bounded operators from $L^2(\Lambda_0)$ to $L^2(\Lambda)$ and $H^1(\Lambda)$ to $L^2(\Lambda_0)$, respectively. On the other hand, the operator S_{12} has a Lipschitz integral kernel

$$\Lambda \times \Lambda_0 \ni (x, y) \mapsto \Phi(x - y)$$

(as the diagonal singularity is excluded), so it defines a compact operator from $L^2(\Lambda_0)$ to $H^1(\Lambda)$ by the subsequent Lemma 12. This shows that $S_{12} N_{21} : L^2(\Lambda) \rightarrow H^1(\Lambda)$ is compact. \square

Lemma 12. *Let $\Lambda, \Lambda_0 \subset \mathbb{R}^n$ be two compact disjoint Lipschitz hypersurfaces and $F : \Lambda \times \Lambda_0 \rightarrow \mathbb{C}$ be a Lipschitz function. Then the operator $S : L^2(\Lambda_0) \rightarrow L^2(\Lambda)$ given by*

$$Sg(x) := \int_{\Lambda_0} F(x, y)g(y) \, ds(y)$$

is actually a compact operator from $L^2(\Lambda_0)$ to $H^1(\Lambda)$

Proof. Let $\{(U_j, V_j, \psi_j)\}_{j \in \{1, \dots, N\}}$, $N \in \mathbb{N}$, be an atlas on Λ consisting of local charts $\psi_j : \mathbb{R}^{n-1} \supset U_j \rightarrow V_j \subset \Lambda$. Let $\chi_j \in C_c^\infty(V_j)$ form a subordinated partition of unity, i.e., $\sum_{j=1}^N \chi_j = 1$ on Λ . Then an equivalent norm on $H^1(\Lambda)$ is given by

$$H^1(\Lambda) \ni g \mapsto \sum_{j=1}^N \|(\chi_j g) \circ \psi_j\|_{H_0^1(U_j)}.$$

Representing

$$(Sg)(\psi_j(u)) = \sum_{j=1}^N \int_{\Lambda_0} F_j(u, y) g(y) \, ds(y),$$

$$F_j(u, y) := \chi_j(\psi_j(u)) F(\psi_j(u), y),$$

we conclude that it is sufficient to show that each of the operators S_j ,

$$S_j : L^2(\Lambda_0) \ni g \mapsto \int_{\Lambda_0} F_j(u, y) g(y) \, ds(y) \in L^2(U_j)$$

defines a compact operator from $L^2(\Lambda)$ to $H_0^1(U_j)$.

Let $j \in \{1, \dots, N\}$ be fixed. As F_j is Lipschitz by construction, for each $k \in \{1, \dots, n-1\}$ the function $\partial_{u_k} F_j$ is bounded. In particular, the integral operators

$$G_{j,k} : L^2(\Lambda_0) \ni g \mapsto \int_{\Lambda_0} \partial_{u_k} F_j(u, y) g(y) \, ds(y) \in L^2(U_j)$$

are Hilbert-Schmidt, therefore, compact. The dominated convergence shows the identity $\partial_k S_j = G_{j,k}$.

Now consider the operator

$$M_j : H_0^1(U_j) \ni f \mapsto \nabla f \in L^2(U_j, \mathbb{C}^{n-1}),$$

where ∇ is the Euclidian gradient. The operator M_j is an isomorphism between $H_0^1(U_j)$ and the closed subspace $\text{ran } M_j$ in $L^2(U_j, \mathbb{C}^{n-1})$, so the inverse

$$M_j^{-1} : \text{ran } M_j \rightarrow H_0^1(U_j)$$

is bounded. We now consider the operator

$$G_j : L^2(\Lambda_0) \ni g \mapsto \begin{pmatrix} G_{j,1} \\ \dots \\ G_{j,n-1} \end{pmatrix} \equiv \begin{pmatrix} \partial_1 S_j \\ \dots \\ \partial_{n-1} S_j \end{pmatrix} \in \text{ran } M_j,$$

which is compact by the above considerations, then

$$S_j = M_j^{-1} G_j : L^2(\Lambda) \rightarrow H_0^1(U_j)$$

is a compact operator being a composition of a compact operator with a bounded operator. \square

3 Transmission problem for Laplacians

Let us consider the geometric configuration described in the introduction, see Fig. 1. Denote by ν the unit normal vector on Σ , exterior with respect to Ω_- , and by ν_0 the exterior unit normal on $\partial\Omega$. Consider the Dirichlet traces

$$\gamma_D^\pm : H^1(\Omega_\pm) \rightarrow L^2(\Sigma), \quad \gamma_D^\partial : H^1(\Omega_+) \rightarrow L^2(\partial\Omega)$$

defined by

$$\gamma_D^\pm f := (f_\pm)|_\Sigma \quad \text{and} \quad \gamma_D^\partial f := (f_+)|_{\partial\Omega} \quad \text{for } f \in H^1(\Omega).$$

Similarly, we let

$$\gamma_N^\pm : H_\Delta^{\frac{3}{2}}(\Omega_\pm) \rightarrow L^2(\Sigma),$$

be the Neumann traces defined by

$$\gamma_N^\pm f := \mp \langle \nu, (\nabla f_\pm)|_\Sigma \rangle \quad \text{for } f_\pm \in H_\Delta^{\frac{3}{2}}(\Omega_\pm).$$

In addition, let $\mu \in \mathbb{R} \setminus \{0, 1\}$. Consider the following linear operator A in $L^2(\Omega)$:

$$\begin{aligned} \text{dom } A &= \{f = (f_+, f_-) \in H_\Delta^{\frac{3}{2}}(\Omega_+) \oplus H_\Delta^{\frac{3}{2}}(\Omega_-) \simeq H_\Delta^{\frac{3}{2}}(\Omega \setminus \Sigma) : \\ &\quad \gamma_D^\partial f_+ = 0 \text{ on } \partial\Omega, \quad \gamma_D^- f_- = \gamma_D^+ f_+ \text{ on } \Sigma, \\ &\quad \mu \gamma_N^- f_- + \gamma_N^+ f_+ = 0 \text{ on } \Sigma\}, \\ A &: (f_+, f_-) \mapsto (-\Delta f_+, -\mu \Delta f_-). \end{aligned}$$

one has obviously $C_c^\infty(\Omega \setminus \Sigma) \subset \text{dom } A$, so A is densely defined. A direct application of the Green formula (Proposition 2) shows that A is a symmetric operator in $L^2(\Omega)$. We are going to find several conditions guaranteeing its self-adjointness. Remark that for $\mu = 1$ the operator A is well-defined as well, but coincides with the Dirichlet Laplacian in Ω , so we exclude this trivial case from the very beginning. We emphasize on the fact that no assumption on the sign of μ is made.

Let $-\Delta_\pm$ be the Dirichlet Laplacian in $L^2(\Omega_\pm)$ and denote

$$B := (-\Delta_+) \oplus (-\mu \Delta_-),$$

which is a self-adjoint operator with compact resolvent in $L^2(\Omega)$. Remark that both A and B are restrictions of the following linear operator T in

$L^2(\Omega)$:

$$\begin{aligned} \text{dom } T &= \{f = (f_+, f_-) \in H_{\Delta}^{\frac{3}{2}}(\Omega_+) \oplus H_{\Delta}^{\frac{3}{2}}(\Omega_-) \simeq H_{\Delta}^{\frac{3}{2}}(\Omega \setminus \Sigma) : \\ &\quad \gamma_D^{\partial} f_+ = 0 \text{ on } \partial\Omega, \gamma_D^- f_- = \gamma_D^+ f_+ \text{ on } \Sigma\}, \\ T &: (f_+, f_-) \mapsto (-\Delta f_+, -\mu \Delta f_-). \end{aligned}$$

Consider the symmetric linear operator Θ in $L^2(\Sigma)$,

$$\text{dom } \Theta = H^1(\Sigma), \quad \Theta = \tilde{N}_0^+ + \mu N_0^-, \quad \tilde{N}_0^+ : H^1(\Sigma) \ni \varphi \mapsto N_0^+(\varphi, 0)|_{\Sigma}.$$

and the linear maps $\Gamma_0, \Gamma_1 : \text{dom } T \rightarrow L^2(\Sigma)$ given by

$$\Gamma_0 f := \gamma_D^- f_- \equiv \gamma_D^+ f_+, \quad \Gamma_1 f := -(\mu \gamma_N^- f_- + \gamma_N^+ f_+) + \Theta \Gamma_0 f.$$

Due to the results of Section 2 we have

$$\text{ran } \Gamma_0 = H^1(\Sigma), \quad \text{ran } \Gamma_1 \subset L^2(\Sigma).$$

For any $f, g \in \text{dom } T$ one has, using the integration by parts,

$$\langle Tf, g \rangle_{L^2(\Omega)} - \langle f, Tg \rangle_{L^2(\Omega)} = \langle \Gamma_1 f, \Gamma_0 g \rangle_{L^2(\Sigma)} - \langle \Gamma_0 f, \Gamma_1 g \rangle_{L^2(\Sigma)}.$$

In addition, the domains of A and B can be represented as

$$\text{dom } A = \{f \in \text{dom } T : \Gamma_0 f \in \text{dom } \Theta, \Gamma_1 f = \Theta \Gamma_0 f\}, \quad \text{dom } B = \ker \Gamma_0.$$

The above properties of T, Γ_0, Γ_1 ensure that the triple $(L^2(\Sigma), \Gamma_0, \Gamma_1)$ is a so-called quasi boundary triple for the adjoint S^* of the symmetric operator $S := T_{\ker \Gamma_0 \cap \ker \Gamma_1}$, see [5, Theorem 2.3], which allows to employ a number of machineries. We are not going to cover the approach in detail (as this would lead to an explosion of the length of the paper and we refer to [6, 18, 19] for a review of various classes of boundary triples), but we remark that the main idea of the approach is to link various properties of A with properties of the “boundary parameter” Θ . One of the main results of the theory is a formula relating the resolvents of A and B , which we are going to state.

Let $z \in \mathbb{C} \setminus \sigma(B)$, then we have a direct sum decomposition

$$\text{dom } T = \text{dom } B \dot{+} \ker(T - z),$$

which shows that $\Gamma_0 : \ker(T - z) \rightarrow \text{ran } \Gamma_0$ is invertible, and we denote the inverse by P_z . In fact, for any $\varphi \in H^1(\Sigma)$ the function $f \equiv (f_+, f_-) := P_z \varphi$ is the solution to

$$\begin{aligned} (-\Delta - z)f_+ &= 0 \text{ in } \Omega_+, & (-\mu \Delta - z)f_- &= 0 \text{ in } \Omega_-, \\ \gamma_D^{\partial} f_+ &= 0, & \gamma_D^+ f_+ &= \varphi = \gamma_D^- f_-. \end{aligned}$$

Due to the results of Section 2 the solution takes the form

$$f_+ := P_z^+(\varphi, 0), \quad f_- := P_{\bar{z}}^- \varphi,$$

where P_z^\pm are the Poisson operators for Ω_\pm and z and uses the natural identifications

$$L^2(\partial\Omega_+) \simeq L^2(\Sigma) \oplus L^2(\partial\Omega), \quad H^1(\partial\Omega_+) \simeq H^1(\Sigma) \oplus H^1(\partial\Omega) \quad \text{etc.}$$

We will also be interested in the operator (Weyl function)

$$M_z := \Gamma_1 P_z : H^1(\Sigma) \rightarrow L^2(\Sigma),$$

which takes the form

$$M_z = \Theta - (\tilde{N}_z^+ + \mu N_{\bar{z}}^-) \equiv (\tilde{N}_0^+ - \tilde{N}_z^+) + \mu(N_0^- - N_{\bar{z}}^-),$$

$$\tilde{N}_z^+ : H^1(\Sigma) \ni \varphi \mapsto N_z^+(\varphi, 0)|_\Sigma,$$

and is compact by Proposition 7.

While P_z is initially defined as an operator from $H^1(\Sigma)$ to $\text{dom } T$, in fact it turns out to be bounded as an operator from $L^2(\Sigma)$ to $L^2(\Omega)$, so the adjoint operator $P_z^* : L^2(\Omega) \rightarrow L^2(\Sigma)$ is well-defined and bounded [5, Prop. 2.6]. The following assertions follow from abstract results on quasi boundary triples [5, Theorem 2.8]:

Proposition 13. (a) For any $z \in \mathbb{C} \setminus \sigma(B)$ one has the equality $\ker(A - z) = P_z(\ker(\Theta - M_z))$. In particular, $\Theta - M_z$ is injective for all $z \in \mathbb{C} \setminus \mathbb{R}$, as the operator A is symmetric.

(b) Let $z \in \mathbb{C} \setminus \sigma(B)$ such that $\Theta - M_z$ is injective and let $f \in L^2(\Omega)$ such that $P_{\bar{z}}^* f \in \text{ran}(\Theta - M_z)$. Then $f \in \text{ran}(A - z)$ and

$$(A - z)^{-1} f = (B - z)^{-1} f + P_z(\Theta - M_z)^{-1} P_{\bar{z}}^* f.$$

Corollary 14. If $\text{ran}(\Theta - M_z) = L^2(\Sigma)$ for all $z \in \mathbb{C} \setminus \mathbb{R}$, then A is self-adjoint.

Proof. By Proposition 13, the assumption ensures $\text{ran}(A - z) = L^2(\Omega)$ for all $z \in \mathbb{C} \setminus \mathbb{R}$. As A is symmetric, this shows the self-adjointness. \square

4 Sufficient conditions for the self-adjointness

Our main tool to verify the assumption of Corollary 14 is:

Theorem 15. *Let $\mu \in \mathbb{R} \setminus \{0, 1\}$ such that the operator*

$$K_\Sigma - \frac{\mu + 1}{2(\mu - 1)} : L^2(\Sigma) \rightarrow L^2(\Sigma)$$

is Fredholm of index m , and let $z \in \mathbb{C} \setminus \mathbb{R}$. Then $\text{ran}(\Theta - M_z)$ is closed with $\dim \text{ran}(\Theta - M_z)^\perp = m$.

Proof. Let $z \in \mathbb{C} \setminus \mathbb{R}$. Remark first that

$$\Theta - M_z \equiv \tilde{N}_z^+ + \mu N_{\bar{z}}^- \equiv \tilde{N}_0^+ + \mu N_0^- + B_1,$$

where $B_1 : H^1(\Sigma) \rightarrow L^2(\Sigma)$ is a compact operator, see Propositions 7 and 10, and

$$(\Theta - M_z)S_\Sigma = \tilde{N}_0^+ S_\Sigma + \mu N_0^- S_\Sigma + B_2, \quad (6)$$

where $B_2 : L^2(\Sigma) \rightarrow L^2(\Sigma)$ is a compact operator. Now consider separately two cases.

Case 1: The operator S_Σ is injective. Then $S_\Sigma : L^2(\Sigma) \rightarrow H^1(\Sigma)$ is bijective by Proposition 3, and then $S_\Sigma^{-1} : H^1(\Sigma) \rightarrow L^2(\Sigma)$ is well defined and bounded. By Proposition 8 we have

$$N_0^- = S_\Sigma^{-1} \left(\frac{1}{2} - K_\Sigma \right)$$

and by Proposition 11 there holds

$$\tilde{N}_0^+ = S_\Sigma^{-1} \left(\frac{1}{2} + K_\Sigma \right) + B_3,$$

where $B_3 : H^1(\Sigma) \rightarrow L^2(\Sigma)$ is a compact operator (the minus sign in front of K_Σ is due to the fact that the normal ν is interior with respect to Ω_+). The substitution of these expressions into (6) gives

$$(\Theta - M_z)S_\Sigma = S_\Sigma^{-1} \left(\frac{1}{2} + K_\Sigma + \frac{\mu}{2} - \mu K_\Sigma \right) S_\Sigma + B_4,$$

where $B_4 := B_2 + B_3 S_\Sigma : L^2(\Sigma) \rightarrow L^2(\Sigma)$ is a compact operator. This can be rewritten as

$$(\Theta - M_z)S_\Sigma = (1 - \mu) S_\Sigma^{-1} \left(K_\Sigma - \frac{\mu + 1}{2(\mu - 1)} \right) S_\Sigma + B_4.$$

Using the symmetrization formula (5) we rewrite the above as

$$(\Theta - M_z)S_\Sigma = (1 - \mu) \left(K_\Sigma^* - \frac{\mu + 1}{2(\mu - 1)} \right) + B_4.$$

Let $K_\Sigma - \frac{\mu+1}{2(\mu-1)}$ be Fredholm of index m , then $K_\Sigma^* - \frac{\mu+1}{2(\mu-1)}$ is Fredholm of index $(-m)$. Due to the compactness of B_4 , the right-hand side is a Fredholm operator of index $(-m)$; in particular, it has a closed range. Then the same applies to the left-hand side. The operator on the left-hand side is injective due to Proposition 13, and due to

$$\text{ran}(\Theta - M_z)S_\Sigma = \text{ran}(\Theta - M(z))$$

the operator $\Theta - M(z)$ has closed range whose codimension is m .

Case 2: The operator S_Σ is not injective. By Proposition 3 the subspace $\ker S$ is one-dimensional, $\ker S = \mathbb{C}f_0$. Define

$$L_\perp^2(\Sigma) := \left\{ f \in L^2(\Sigma) : \int_\Sigma f \bar{f}_0 ds = 0 \right\}, \quad H_\perp^1(\Sigma) := H^1(\Sigma) \cap L_\perp^2(\Sigma),$$

then the operator

$$S_0 := \left(S_\Sigma : L_\perp^2(\Sigma) \rightarrow H_\perp^1(\Sigma) \right)$$

is an isomorphism. Remark that $H_\perp^1(\Sigma)$ is a closed subspace of $H^1(\Sigma)$ of codimension one. Let P be the orthogonal projection from $L^2(\Sigma)$ onto $L_\perp^2(\Sigma)$. Then, from Propositions 8 and 11 one has

$$N_0^- = S_0^{-1}P\left(\frac{1}{2} - K_\Sigma\right) \quad \text{and} \quad \tilde{N}_0^+ = S_0^{-1}\left(\frac{1}{2} + K_\Sigma\right) + \tilde{B}_3$$

with $\tilde{B}_3 : H^1(\Sigma) \rightarrow L^2(\Sigma)$ a compact operator. Here we used the fact that $1 - P$ is an operator with a rank equal to 1, and hence a compact operator. Now, using that $S_0^{-1}S_\Sigma = P$ and $PS_\Sigma = S_\Sigma$, the same arguments as before yield

$$\begin{aligned} (\Theta - M(z))S_\Sigma &= S_0^{-1}\left(P\left(\frac{1}{2} + K_\Sigma\right) + \frac{\mu}{2} - \mu K_\Sigma\right)S_\Sigma + \tilde{B}_4, \\ &= S_0^{-1}\left(\frac{1+\mu}{2} + PSK_\Sigma^* - \mu S_\Sigma K_\Sigma^*\right) + B_5, \\ &= (1-\mu)\left(K_\Sigma^* - \frac{\mu+1}{2(\mu-1)}\right) + B_6, \end{aligned}$$

where \tilde{B}_4 , B_5 and B_6 are compact operators from $L^2(\Sigma)$ into itself. As above we deduce that $(\Theta - M(z))S_\Sigma$ is Fredholm with index $-m$ which entails that $(\Theta - M(z))S_\Sigma$ has a closed range. Then, Proposition 3 yields that $\text{ran}(\Theta - M(z)) = \text{ran}(\Theta - M(z))S_\Sigma + V$ where V is a subspace of dimension 1. Since $H_\perp^1(\Sigma)$ is a closed subspace of $H^1(\Sigma)$ it follows that $\text{ran}(\Theta - M(z))$ is closed. From this and the fact that S_Σ is Fredholm with index zero, we then deduce from Proposition 13 that $\dim \text{ran}(\Theta - M(z))^\perp = m$ and this proves the claim. \square

For what follows, for a bounded linear operator C in a Banach space it will be convenient to denote

$$\begin{aligned}\sigma_{\text{ess}}(C) &:= \{z \in \mathbb{C} : C - z \text{ is not a Fredholm operator}\}, \\ \sigma_{\text{ess}}^0(C) &:= \{z \in \mathbb{C} : C - z \text{ is not a zero index Fredholm operator}\}.\end{aligned}$$

One has the trivial inclusions $\sigma_{\text{ess}}(C) \subset \sigma_{\text{ess}}^0(C) \subset \sigma(C)$ and the equalities

$$\sigma(C^*) = \overline{\sigma(C)}. \quad \sigma_{\text{ess}}(C^*) = \overline{\sigma_{\text{ess}}(C)}, \quad \sigma_{\text{ess}}^0(C^*) = \overline{\sigma_{\text{ess}}^0(C)}.$$

In addition define the spectral radius $r(C)$ and the essential spectral radius $r_{\text{ess}}(C)$ by

$$r(C) := \sup \{|\lambda| : \lambda \in \sigma(C)\}, \quad r_{\text{ess}}(C) := \sup \{|\lambda| : \lambda \in \sigma_{\text{ess}}(C)\};$$

one has clearly

$$r_{\text{ess}}(C) \leq r(C), \quad r(C^*) = r(C), \quad r_{\text{ess}}(C^*) = r_{\text{ess}}(C).$$

Recall that the index of $C - \lambda$ is constant for λ in each connected component of $C \setminus \sigma_{\text{ess}}(C)$. As the index is zero for $|\lambda| > \|C\|$, we conclude that

$$\sigma_{\text{ess}}(C) \subset \{\lambda \in C : |\lambda| \leq r_{\text{ess}}(C)\}.$$

With these conventions and observations, Theorem 15 has the following direct consequence.

Theorem 16. *Let $\mu \in \mathbb{R} \setminus \{0, 1\}$ such that*

$$\frac{\mu + 1}{2(\mu - 1)} \notin \sigma_{\text{ess}}^0(K_\Sigma), \tag{7}$$

then A is self-adjoint with compact resolvent. The condition (7) is satisfied, in particular, if

$$\left| \frac{\mu + 1}{2(\mu - 1)} \right| > r_{\text{ess}}(K_\Sigma).$$

Proof. We only prove that $(A - z)^{-1}$ is compact. Since $\Theta : H^1(\Sigma) \rightarrow L^2(\Sigma)$ is bounded, by the closed graph theorem it follows that $(\Theta - M(z))^{-1}$ is well-defined and bounded from $L^2(\Sigma)$ to $H^1(\Sigma)$, and hence a compact operator in $L^2(\Sigma)$ due to the compactness of the embedding $H^1(\Sigma) \hookrightarrow L^2(\Sigma)$. Since P_z is bounded from $L^2(\Sigma)$ to $L^2(\Omega)$, from the resolvent formula in Proposition 13 we deduce that $(A - z)^{-1}$ is compact. \square

We now establish several concrete situations in which (7) is satisfied. First remark that [20, Theorem 2]

$$r(K_\Sigma) \equiv r(K_\Sigma^*) \leq \frac{1}{2}. \tag{8}$$

Theorem 17. *The operator A is self-adjoint for any $\mu > 0$ without any additional assumption on Σ .*

Proof. Let $\mu > 0$ with $\mu \neq 1$, then due to (8) we have

$$\left| \frac{\mu + 1}{2(\mu - 1)} \right| > \frac{1}{2} \geq r(K_\Sigma) \geq r_{\text{ess}}(K_\Sigma),$$

and the claim follows by Corollary 16. The case $\mu = 1$ is trivial. \square

Recall [24] that $\text{VMO}(\Sigma)$ stands for the space of functions of vanishing means oscillation on Σ . The bounded Lipschitz domains with normals in VMO are those domains "without corners". This includes all bounded C^1 -smooth domains.

Theorem 18. *Let Σ be such that $\nu \in \text{VMO}(\Sigma)$, which is satisfied, in particular for C^1 -smooth Σ . Then the operator A is self-adjoint for any $\mu \in \mathbb{R} \setminus \{-1, 0\}$.*

Proof. The operator K_Σ is compact on $L^2(\Sigma)$ by [24, Theorem 4.47], so $\sigma_{\text{ess}}^0(K_\Sigma) = \{0\}$ and $r_{\text{ess}}(K_\Sigma^*) = 0$, and for $\mu \notin \{-1, 0, 1\}$ the claim follows by Corollary 16. The case $\mu = 1$ is trivially covered. \square

Remark 19. So far we have no self-adjointness condition for the interesting case $\mu < 0$ and Σ with corners. In fact, if for a given Σ one can improve (8) to

$$r := r_{\text{ess}}(K_\Sigma) < \frac{1}{2},$$

then for $\mu \notin \{0, 1\}$ there holds

$$\left| \frac{\mu + 1}{2(\mu - 1)} \right| > r \text{ if and only if } \mu \notin I_r := \left[-\frac{1 + 2r}{1 - 2r}, -\frac{1 - 2r}{1 + 2r} \right],$$

i.e. the self-adjointness of the operator A is also guaranteed for all negative μ outside the "critical interval" I_r . At the same time, the inequalities

$$r(K_\Sigma) < \frac{1}{2}, \quad r_{\text{ess}}(K_\Sigma) < \frac{1}{2}$$

represent central conjectures in the theorem of Neumann-Poincaré operators, which are still unsolved in the general form. The papers [15, 45] contain a rather complete bibliography and a review of available results on the spectral radii. Let us discuss an important particular case in greater detail.




Figure 2: Plots of $\omega \mapsto a(\omega)$ (solid line) and $\omega \mapsto b(\omega)$ (dashed line).

Theorem 20. *Let $n = 2$ and Σ be a curvilinear polygon with C^1 -smooth edges and with N interior angles $\omega_1, \dots, \omega_N \in (0, 2\pi) \setminus \{0\}$. Let $\omega \in (0, \pi)$ be the sharpest angle, i.e.*

$$\frac{|\pi - \omega|}{2} = \max_k \frac{|\pi - \omega_k|}{2},$$

then the operator A is self-adjoint for all $\mu \neq 0$ with

$$\mu \notin \left[-\frac{1}{a(\omega)}, -a(\omega) \right] \text{ for } a(\omega) := \tan^2 \frac{\omega}{4} \equiv \frac{1 - \cos \frac{\omega}{2}}{1 + \cos \frac{\omega}{2}}. \quad (9)$$

Proof. By [41, Theorem 6] there holds

$$r_{\text{ess}}(K_\Sigma) = \frac{1}{2} \max_k \sin \frac{|\pi - \omega_k|}{2},$$

and the rest follows from the computations in Remark 19 with the help of elementary trigonometric identities. \square

Remark 21. The case of curvilinear polygons was already considered (with some non-essential restrictions) in [9, 12, 17]. These works started with a lower regularity, i.e. they considered the operator A_0 defined in the same way as A but allowing for $u_\pm \in H_\Delta^1(\Omega_\pm)$ instead of $u_\pm \in H_\Delta^{\frac{3}{2}}(\Omega_\pm)$ (with a suitably adapted definition of boundary traces), and it was shown that A_0 is self-adjoint for all $\mu \neq 0$ such that

$$\mu \notin \left[-\frac{1}{b(\omega)}, -b(\omega) \right] \text{ for } b(\omega) := \frac{\omega}{2\pi - \omega}. \quad (10)$$

A simple analysis shows that the critical interval in our bound (9) is larger than the one in (10), which is the price to pay to have the higher $H^{\frac{3}{2}}$ -regularity. A numerical comparison of the two critical intervals is shown in Figure 2.

Remark 22. We are not aware of efficient estimates for $r_{\text{ess}}(K_\Sigma)$ if $n \geq 3$ and Σ has corners. In [28, Section 4] it was shown that if Σ is an arbitrary three-dimensional polyhedron, then $r_{\text{ess}}(K_{\text{ess}}) < \frac{1}{2}$, but no constructive upper bound was available. In [23, Theorem 5.5] one finds an expression for the $\sigma_{\text{ess}}(K_{\text{ess}})$ for the case of rotationally invariant conical singularities in three dimensions in terms of modified Legendre functions, but the expressions obtained did not contain an explicit bound of $r_{\text{ess}}(K)$.

Remark 23. We believe that a more general statement should be valid: if $K - \frac{\mu+1}{2(\mu-1)}$ is Fredholm of index m , then A is a closed symmetric operator with deficiency indices (m, m) . This should follow by a suitable extension of the resolvent formula, but which would in turn require some non-evident and adapted extensions of the Dirichlet/Neumann traces. We prefer to leave this question for future work.

Acknowledgments

The authors were in part supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – 491606144.

References

- [1] K. Ando, H. Kang, Y. Miyanishi, M. Putinar: *Spectral analysis of Neumann-Poincaré operator*. Rev. Roumaine Math. Pures. Appl. **66**:3-4 (2021) 545–575.
- [2] J. Behrndt, F. Gesztesy, M. Mitrea: *Sharp boundary trace theory and Schrödinger operators on bounded Lipschitz domains*. Mem. Amer. Math. Soc. (in press). Preprint arXiv:2209.09230.
- [3] J. Behrndt, M. Holzmann, C. Stelzer, G. Stenzel: *Boundary triples and Weyl functions for Dirac operators with singular interactions*. Rev. Math. Phys. (in press). Preprint arXiv:2211.05191.
- [4] J. Behrndt, D. Krejčířík: *An indefinite Laplacian on a rectangle*. J. Anal. Math. **134** (2018) 501-522.
- [5] J. Behrndt, M. Langer: *Boundary value problems for elliptic partial differential operators on bounded domains*. J. Funct. Anal. **243** (2007) 536–565.
- [6] J. Behrndt, M. Langer: *Elliptic operators, Dirichlet-to-Neumann maps and quasi boundary triples*. In S. Hassi, H. de Snoo, F. H. Szafraniec (Eds.): *Operator methods for boundary value problems* (London Math. Soc. Lecture Note Series, Vol. 404, Cambridge University Press, 2012) 121–160.
- [7] J. Behrndt, M. Holzmann, T. Ourmières-Bonafos, K. Pankrashkin: *Two-dimensional Dirac operators with singular interactions supported on closed curves*. J. Funct. Anal. **279** (2020) 108700.

- [8] B. Benhellal, K. Pankrashkin: *Curvature contribution to the essential spectrum of Dirac operators with critical shell interactions*. Pure Appl. Anal. (in press). Preprint arXiv:2211.10264.
- [9] A.-S. Bonnet-Ben Dhia, L. Chesnel, P. Ciarlet Jr.: *T-coercivity for scalar interface problems between dielectrics and metamaterials*. ESAIM: M2AN **46** (2012) 1363–1387.
- [10] A.-S. Bonnet-Ben Dhia, L. Chesnel, X. Claeys: *Radiation condition for a non-smooth interface between a dielectric and a metamaterial*. Math. Models Meth. App. Sci. **23** (2013) 1629–1662.
- [11] A.-S. Bonnet-Ben Dhia, L. Chesnel, M. Rihani: *Maxwell’s equations with hypersingularities at a conical plasmonic tip*. J. Math. Pures Appl. **161** (2022) 70–110.
- [12] A.-S. Bonnet-Ben Dhia, M. Dauge, K. Ramdani: *Analyse spectrale et singularités d’un problème de transmission non-coercive*. C. R. Acad. Sci. Paris **328** (1999) 717–720.
- [13] G. Bouchitté, B. Schweizer: *Cloaking of small objects by anomalous localized resonance*. Quart. J. Mech. Appl. Math. **63** (2010) 438–463.
- [14] C. Cacciapuoti, K. Pankrashkin, A. Posilicano: *Self-adjoint indefinite Laplacians*. J. Anal. Math. **139** (2018) 155–177.
- [15] S. N. Chandler-Wilde, R. Hagger, K.-M. Perfekt, J. A. Virtanen: *On the spectrum of the double-layer operator on locally-dilation-invariant Lipschitz domains*. Num. Math. **153** 635–699.
- [16] M. Costabel, E. Stephan: *A direct boundary intergal equation method for transmission problems*. J. Math. Anal. Appl. **106** (1985) 367–413.
- [17] M. Dauge, B. Texier: *Problèmes de transmission non coercifs dans des polygones*. Report IRMAR 97-27 (Rennes, 1997), arXiv:1102.1409.
- [18] V. Derkach, S. Hassi, M. Malamud: *Generalized boundary triples, I. Some classes of isometric and unitary boundary pairs and realization problems for subclasses of Nevanlinna functions*. Math. Nachr. **293** (2020) 1278–1327.
- [19] V. Derkach, S. Hassi, M. Malamud: *Generalized boundary triples, II. Some applications of generalized boundary triples and form domain invariant Nevanlinna functions*. Math. Nachr. **295** (2022) 1113–1162.
- [20] L. Escauriaza, E. B. Fabes, G. Verchota: *On a regularity theorem for weak solutions to transmission problems with internal Lipschitz boundaries*. Proc. Amer. Math. Soc. **115** (1992) 1069–1076.
- [21] D. Grieser: *The plasmonic eigenvalue problem*. Rev. Math. Phys. **26** (2014) 1450005.
- [22] L. Grubišić, V. Kostykin, K. A. Makarov, K. Veselić: *Representation theorems for indefinite quadratic forms revisited*. Mathematika **59** (2013) 169–189.
- [23] J. Helsing, K.-M. Perfekt: *The spectra of harmonic layer potential operators on domains with rotationally symmetric conical points*. J. Math. Pures Appl. **118** (2018) 235–287.

- [24] S. Hofmann, M. Mitrea, M. Taylor: *Singular integrals and elliptic boundary problems on regular Semmes-Kenig-Toro domains*. Int. Math. Res. Not. **2010** (2010) 2567–2865.
- [25] H. Kettunen, M. Lassas, P. Ola: *On absence and existence of the anomalous localized resonance without the quasi-static approximation*. SIAM J. Appl. Math. **78** (2018) 609–628.
- [26] R. V. Kohn, J. Lu, B. Schweizer, M. I. Weinstein: *A variational perspective on cloaking by anomalous localized resonance*. Comm. Math. Phys. **328** (2014) 1–27.
- [27] V. Kostrykin: *The $\operatorname{div}(A \operatorname{grad})$ operator without ellipticity. Self-adjointness and spectrum*. In M. van den Berg, D. Grieser, T. Hoffmann-Ostenhof, I. Polterovich (Eds.), *Geometric aspects of spectral theory*. Oberwolfach Reports **9** (2012) 2061–2063.
- [28] M. de León-Contreras, K.-M. Perfekt: *The quasi-static plasmonic problem for polyhedra*. Math. Ann. **387** (2023) 1533–1577.
- [29] W. McLean: *Strongly elliptic systems and boundary integral equations*. Cambridge University Press, 2000.
- [30] D. Medková: *The Laplace equation*. Springer, 2018.
- [31] Y. Miyanishi: *Weyl’s law for the eigenvalues of the Neumann-Poincaré operators in three dimensions: Willmore energy and surface geometry*. Adv. Math. **406** (2022) 108547.
- [32] Y. Miyanishi, G. Rozenblum: *Eigenvalues of the Neumann-Poincaré operator in dimension 3: Weyl’s law and geometry*. St. Petersburg Math. J. **31** (2020) 371–386.
- [33] H. Mohsen, S. Labrunie, V. Nistor: *Estimations polynomiales pour les problèmes de transmission sur des domaines à bords plats*. Tunis. J. Math. **4** (2022) 755–777.
- [34] H. Mohsen, S. Labrunie, V. Nistor: *Polynomial bounds for the solutions of parametric transmission problems on smooth, bounded domains*. Preprint arXiv:2308.06215.
- [35] H.-M. Nguyen: *Asymptotic behavior of solutions to the Helmholtz equations with sign changing coefficients*. Tran. Amer. Math. Soc. **367** (2015) 6581–6595.
- [36] H.-M. Nguyen: *Limiting absorption principle and well-posedness for the Helmholtz equation with sign changing coefficients*. J. Math. Pures Appl. **106** (2016) 342–374.
- [37] P. Ola: *Remarks on a transmission problem*. J. Math. Anal. Appl. **196** (1995) 639–658.
- [38] K. Pankrashkin: *On self-adjoint realizations of sign-indefinite Laplacians*. Rev. Roumaine Math. Pures Appl. **64:2-3** (2019) 345–372.
- [39] S. Schmitz: *Representation theorems for indefinite quadratic forms without spectral gap*. Integr. Equ. Oper. Th. **83** (2015) 73–94.

- [40] S. Schmitz: *Representation theorems for indefinite quadratic forms and applications*. Dissertation, Mainz, 2014. Full text available at <http://nbn-resolving.org/urn:nbn:de:hebis:77-37267>.
- [41] V. Yu. Shelepov: *On the index and spectrum of integral operators of potential type along Radon curves*. Math. USSR Sb. **70** (1991) 175–203.
- [42] G. Verchota: *Layer potentials and regularity for the Dirichlet problem for Laplace's equation in Lipschitz domains*. J. Funct. Anal. **59** (1984) 572–611.
- [43] V. G. Veselago: *The electrodynamics of substances with simultaneously negative values of ε and μ* . Sov. Phys. Usp. **10** (1968) 509–514.
- [44] M. Waurick: *Homogenisation of laminated metamaterials and the inner spectrum*. Preprint arXiv:2210.04650.
- [45] W. L. Wendland: *On the double layer potential*. In A. Cialdea, F. Lanzara, P. E. Ricci (Eds.): *Analysis, partial differential equations and applications. The Vladimir Maz'ya anniversary volume*. (Oper. Theory Adv. Appl., Vol. 193, Birkhäuser, 2009) 319–334.