

# SECTION 4. ACKNOWLEDGEMENTS, REFERENCES & APPENDICES

## 1. ACKNOWLEDGEMENTS

This is a contribution to the LEDA Traitbase project. The LEDA Traitbase project is funded through the 5th framework of the EC under the Energy, Environment, and Sustainable Development programme (EESD), contract number EVR1-CT-2002-40022.

## 2. REFERENCES

### 2.1. LITERATURE REFERENCES

- Aksoy, A., Dixon, J.M. & Hale, W.H.G. (1998) Biological Flora of the British Isles: *Capsella bursa-pastoris* (L.) Medikus (*Thlaspi bursa-pastoris* L., *Bursa bursa-pastoris* (L.) Shull, *Bursa pastoris* (L.) Weber). *Journal of Ecology*, 86, 171-186.
- Arber, A. (1920) Water plants. A study of aquatic angiosperms. Cambridge University Press.
- Askw, A.P., Corker, D., Hodkinson, D.J. & Thompson, K. (1996) A new apparatus to measure the rate of fall of seeds. *Functional Ecology*, 11, 121-125.
- Atkinson, M.D. & Atkinson, E. (2002) Biological Flora of the British Isles: *Sambucus nigra* L. *Journal of Ecology*, 90, 895-923.
- Augsburger, C.K. (1986) Morphology and dispersal potential of wind-dispersed diaspores of neotropical trees. *American Journal of Botany*, 73(3), 353-363.
- Bakker, J.P. (1987) Restoration of species-rich grasslands after a period of fertiliser application. In: *Disturbance in Grasslands* (eds J. Van Andel, J.P. Bakker & R.W. Snaydon), pp. 185-200. Junk Publishers, Dordrecht.
- Bakker, J.P. (1989) Nature Management by Grazing and Cutting. Kluwer, Dordrecht.
- Bakker, J.P., Bos, A.F., Hoogveld, J. & Muller, H.J. (1991) The role of the seed bank in restoration management of semi-natural grasslands. In: *Terrestrial and aquatic ecosystems; perturbation and recovery* (ed. O. Ravera), pp. 449-455. Ellis Morwood.
- Bakker, J.P., Elzinga, J.A. & De Vries, Y. (2002) Effects of long-term cutting in a grassland system: perspectives for restoration of plant communities on nutrient-poor soils. *Applied Vegetation Science*, 5, 107-120.
- Bakker, J.P., Poschod, P., Strykstra, R.J., Bekker, R.M. & Thompson, K. (1996) Seed banks and seed dispersal: important topics in restoration ecology. *Acta Botanica Neerlandica*, 45, 461-490.
- Barkman, J.J. (1979) The investigation of vegetation texture and structure. In: *The study of vegetation* (ed. Werger, M.J.A.), pp. 123-160. Junk, The Hague.
- Barkman, J.J. (1988) New Systems of Plant growth forms and phenological plant types. In: *Plant form and vegetation structure* (eds M.J.A. Werger P.J.M Van der Aart, H.J. Doring, J.T.A. Verhoeven), pp. 9-44. S. P. B. Publishers, The Hague.
- Baskin C.C. & Baskin, J.M. (1998) *Seeds - Ecology, Biogeography and Evolution of Dormancy and Germination*. Academic Press, California.
- Begon, M., Mortimer, M. & Thompson, D.J. (1997) *Populationökologie*. Spektrum Akademischer Verlag, Heidelberg.
- Bekker, R.M., Schaminée, J.H.J., Bakker, J.P. & Thompson, K. (1998) Seed bank characteristics of Dutch plant communities. *Acta Botanica Neerlandica*, 47, 15-26.
- Bell, A.D. (1991) *An Illustrated Guide to Flowering Plant Morphology*. Oxford University Press.
- Bellingham, P.J. & Sparrow, A.D. (2000) Resprouting as a life history strategy in woody plant communities. *Oikos*, 89, 409-416.
- Bill, H.-C., Poschod, P., Reich, M. & Plachter, H. (1999) Experiments and observations on seed dispersal by running water in an Alpine floodplain. *Bulletin of the Geobotanical Institute ETH*, 65, 13-28.
- Bill, H.-C. (2000) *Besiedlungsdynamik und Populationsbiologie charakteristischer Pionierpflanzenarten nordalpiner Wildflüsse*. Wissenschaft in Dissertationen Band 557, Görich & Weiershäuser GmbH, Marburg.

- Böcker, R., Kowarik, I. & Bornkamm, R. (1983) Untersuchungen zur Anwendung der Zeigerwerte nach Ellenberg. *Verhandlungen der Gesellschaft für Ökologie*, 11, 35-56.
- Bonn, S. & Poschold, P. (1998) *Ausbreitungsbiologie der Pflanzen Mitteleuropas*. Quelle & Meyer Verlag, UTB, Wiesbaden.
- Bonn, S. (XXXX) Assessment of endozoochorous dispersal potential in plant species - a methodological approach. In prep.
- Brack, L. & Wood, G.B. (1997) *Forest mensuration*. URL: <http://sres.anu.edu.au/associated/mensuration/BrackandWood1998>.
- Burns, I.G. (1991) Short- and long-term effects of a change in the spatial distribution of nitrate in the root zone on N uptake, growth and root development of young lettuce plants. *Plant, Cell & Environment*, 14, 21-33.
- Cain, M.L., Milligan, B.G. & Strand, A.E. (2000) Long-distance seed dispersal in plant populations. *American Journal of Botany*, 87, 1217-1227.
- Caldwell, M.M., Manwaring J.H. & Durham, S.L. (1991a) The microscale distribution of neighbouring plant roots in fertile soil microsites. *Functional Ecology*, 5, 765-772.
- Caldwell, M.M., Manwaring, J.H. & Jackson, R.B. (1991b) Exploitation of phosphate from fertile soil microsites by three Great Basin perennials when in competition. *Functional Ecology*, 5, 757-764.
- Castro-Díez, P., Puyravaud, J.P., Cornelissen, J.H.C. & Villar-Salvador, P. (1998) Stem anatomy and relative growth rate in seedlings of a wide range of woody plant species and types. *Oecologia*, 116, 57-66.
- Cerabolini, B., Ceriani, R., Caccianiga, M., Andreis, R.D. & Raimondi, B. (2003) Seed size, shape and persistence in the soil: a test on Italian flora from Alps to Mediterranean coasts. *Seed Science Research*, 13, 75-85.
- Cerda, A. & Garcia-Fayos, P. (2002) The influence of seed size and shape on their removal by water erosion. *Catena*, 48, 293-302.
- Chapman, D. (ed.) (1996) *Water quality assessments, a guide to the use of biota, sediments and water in environmental monitoring*. Chapman & Hall, London.
- Clausen, P., Nolet, B.A., Fox, A.D. & Klaassen, M. (2002) Long-distance endozoochorous dispersal of submerged macrophyte seeds by migratory waterbirds in Northern Europe – a critical review of possibilities and limitations. *Acta Oecologica*, 23, 191-203.
- Cody, W.J. & Crompton, C.W. (1975) The Biology of Canadian weeds: 15. *Pteridium aquilinum* (L.) Kuhn. *Canadian Journal of Plant Science*, 55, 1059-1072.
- Cole, C.V., Innis, G.S. & Stewart, J.W.B. (1977) Simulation of phosphorus cycling in semi-arid grassland. *Ecology*, 58, 1-15.
- Cook, C.D.K. (1987) Vegetative growth and genetic mobility in some aquatic weeds. Differentiation patterns in higher plants (ed. K.M. Urbanska), pp. 217–225. Academic Press, London.
- Cornelissen, J.H.C., Lavorel, S., Garnier, E., Diaz, S., Buchmann, N., Gurvich, D.E., Reich, P.B., Ter Steege, H., Morgan, H.D., Van der Heijden, M.G.A., Pausas, J.G. and Poorter, H. (2003) Handbook of protocols for standardised and easy measurement of plant functional traits worldwide. *Australian Journal of Botany*, 51, 335-380.
- Crawley, M.J. & Long, C.R. (1995) Alternate bearing, predator satiation and seedling recruitment in *Quercus robur* L. *Journal of Ecology*, 83, 683-696.
- Dalling, J.W. & Hubbell, S.P. (2002) Seed size, growth rate and gap micro-site conditions as determinants of recruitment success for pioneer species. *Journal of Ecology*, 90, 557-568.
- Danvind, M. & Nilsson, C. (1997) Seed floating ability and distribution of alpine plants along a northern Swedish river. *Journal of Vegetation Science*, 8, 271-276.
- De Kroon, H., van Groenendael, J. (eds.) (1997) *The ecology and evolution of clonal plants*. Backhuys Publishers, Leiden.
- De Lyon, M.J.H. & Roelofs, J.G.M. (1986) *Waterplanten in relatie tot waterkwaliteit en bodemgesteldheid, deel 1 & deel 2*. Report of the department of Aquatic Ecology, University of Nijmegen, Nijmegen.
- Del Tredici, P. (2001) Sprouting in temperate trees: a morphological and ecological review. *Botanical Review*, 67, 121-140.
- Den Hartog, C. & Segal, S. (1964) A new classification of the water plant communities. *Acta Botanica Neerlandica*, 13, 367-393.
- Diaz, S., Cabido, M., Zak, M., Martínez-Carretero, E. & Aranibar, J. (1999) Plant functional traits, ecosystem structure, and land-use history along a climatic gradient in central-western Argentina. *Journal of Vegetation Science*, 10, 651-660.
- Diekmann, M. (1995) Use and improvement of Ellenberg's indicator values in deciduous forests of the Boreo-nemoral zone in Sweden. *Ecography* 18, 178-189.
- Diemer, M. & Körner, Ch. (1996) Lifetime leaf carbon balances of herbaceous perennial plants from low and high altitudes in the central Alps. *Functional Ecology*, 10, 33-43.

- Diggle, P.K. (1997) Extreme preformation in alpine *Polygonum viviparum*: An architectural and developmental analysis. *American Journal of Botany* 84, 154-169.
- Domin, K. (1925) Rostlinné tvarosloví. Základy srovnávací a popisné morfologie rostlin. Československá Botanická společnost v Praze.
- Drew, M.C., Saker, L.R. & Ashley, T.W. (1973) Nutrient supply and the growth of the seminal root system in barley. I. The effect of nitrate concentration on the growth of axes and laterals. *Journal of Experimental Botany*, 24, 1189-1202.
- EEA (2002) EUNIS habitat classification. Version 2.3. URL: <http://mrw.wallonie.be/dgrne/sibw/EUNIS/home.html>.
- Ehrlen, J. & Van Groenendael, J.M. (1998) The trade-off between dispersability and longevity – an important aspect of plant species diversity. *Applied Vegetation Science*, 1, 29-36.
- Ellenberg, H., Weber, H.E., Dull, R., Wirth, V., Werner, W. & Paulissen, D. (1991) Zeigerwerte von Pflanzen in Mitteleuropa. *Scripta Geobotanica*, 18, 1-248.
- Enquist, B.J., West, G.B., Charnov, E.L. & Brown, J.H. (1999) Allometric scaling of production and life-history variation in vascular plants. *Nature*, 401, 907-911.
- Eriksson, O. & Fröberg, H. (1996) Windows of opportunity for recruitment in long-lived clonal plants: experimental studies of seedling establishment in *Vaccinium* shrubs. *Canadian Journal of Botany*, 74, 1369-1374.
- Eriksson, O. & Jakobsson, A. (1998) Abundance, distribution and life histories of grassland plants: a comparative study of 81 species. *Journal of Ecology*, 86, 922-933.
- Escarre, J. & Thompson, J.D. (1991) The effects of successional habitat variation and time of flowering on seed production in *Rumex acetosella*. *Journal of Ecology*, 79, 1099-1112.
- EUNIS (2002) *EUNIS Habitat classification glossary*. URL: <http://mrw.wallonie.be/dgrne/sibw/EUNIS/eunis.gloss.html>.
- FAO (1998) World Reference Base for Soil Resources. Food and Agriculture organization of the United Nations, International Society of Soil Science. World Soil Resources Report No 84. Download available at <http://www.fao.org/WAICENT/FAOINFO/AGRICULT/AGL/agll/wrb/wrbdocs.stm>.
- FAO (2001) Lecture Notes on Major Soils of the World.. Food and Agriculture organization of the United Nations, International Society of Soil Science. World Soil Resources Report No 94. Download available at <http://www.fao.org/WAICENT/FAOINFO/AGRICULT/AGL/agll/wrb/wrbdocs.stm>.
- Fenner, M. (1992) *Seeds: the ecology of regeneration in plant communities*. CAB International, Wallingford, United Kingdom.
- Fenner, M. (ed.) (1992) *Seeds: the Ecology of Regeneration in Plant Communities*. CAB International, Wallingford.
- Fermania, T.W. (2000) Controlling turfgrass pests: Weed biology. URL: <http://www.turf.uiuc.edu/teaching/NRES300/weed%20biology.htm>.
- Fernández-Santos, B., Gómez-Gutiérrez, J.M. & Moreno-Marcos, G. (1999) Effects of disturbance caused by traditional Spanish rural land use on the regeneration of *Cytisus multiflorus*. *Applied Vegetation Science*, 2, 239-250.
- Fischer, S.F., Poschlod, P. & Beinlich, B. (1996) Experimental studies on the dispersal of plants and animals on sheep in calcareous grasslands. *Journal of Applied Ecology*, 33, 1206-1222.
- Gardener, C.J., Mclvor, J.G. & Jansen, A. (1993) Passage of legume and grass seeds through the digestive tract of cattle and their survival in faeces. *Journal of Applied Ecology*, 30, 63-74.
- Gardner, G. (1977) The reproductive capacity of *Fraxinus excelsior* on the Derbyshire Limestone. *Journal of Ecology*, 65, 107-118.
- Garnier, E. & Laurent, G. (1994) Leaf anatomy, specific mass and water content in congeneric annual and perennial grass species. *New Phytologist*, 128, 725-736.
- Garnier, E., Cordonnier, P., Guillem, J.-L. & Sonié, L. (1997) Specific leaf area and leaf nitrogen concentration in annual and perennial grass species growing in Mediterranean old-fields. *Oecologia*, 111, 490-498.
- Garnier, E., Shipley, B., Roumet, C. & Laurent, G. (2001) A standardised protocol for the determination of specific leaf area and leaf dry matter content. *Functional Ecology*, 15, 688-695.
- Gaudet, C. L. & Keddy, P.A. (1988) A comparative approach to predicting competitive ability from plant traits. *Nature*, 334, 242-243.
- Golterman, H.L. (ed.) (1969) *Methods for chemical analysis of fresh waters*. International Biological Program Handbook 8. Blackwell, Oxford.
- Goodwin, M.J, Parkinson, R.J., Williams, E.N.D. & Tallowin, J.R.B. (1998) Soil phosphorus extractability and uptake in a *Cirsio moliniatum* fen-meadow and an adjacent *Holcus lanatus* pasture on the culm measures. *Agriculture, Ecosystems and Environment*, 70, 169-179.
- Greene, D.F. & Johnson, E.A. (1989) A model of wind dispersal of winged or plumed seeds. *Ecology*, 70, 339-347.

- Grime, J.P., Hodgson, J.G. & Hunt, R. (1988) *Comparative Plant Ecology*. Unwin Hyman, London.
- Grissino-Mayer, H.D. (1999) *Introduction to Soil Science*. URL: [www.valdosta.edu/~grissino/geol3710/soil14.htm](http://www.valdosta.edu/~grissino/geol3710/soil14.htm)
- Gross, K.L. (1990) A comparison of methods for estimating seed numbers in the soil. *Journal of Ecology*, 78, 1079-1093.
- Groth, A.T., Lovett-Doust, L. & Lovett-Doust, J. (1996) Population density and module demography in *Trapa natans* (Trapaceae), an annual, clonal aquatic macrophyte. *American Journal of Botany*, 83, 1406-1415.
- Hacke, U.G., Sperry, J.S. & Pitterman, J. (2000) Drought experience and cavitation resistance in six shrubs from the Great Basin, Utah. *Basic and Applied Ecology*, 1, 31-41.
- Hacke, U.G., Sperry, J.S., Pockman, W.T., Davis, S.D. & McCulloh, K.A. (2001) Trends in wood density and structure are linked to prevention of xylem implosion by negative pressure. *Oecologia*, 126, 457-461.
- Hamdoun, A.M. (1972) Regenerative capacity of root fragments of *Cirsium arvense* (L.) SCOP. *Weed Research*, 12, 128-136.
- Hanf, M. (1990) *Ackerunkräuter Europas mit ihren Keimlingen und Samen*. 3. Aufl., BLV München, DLG-Verl. Frankfurt/M., Landwirtschaftsverlag Münster-Hiltrup, Österr. Agrarverlag Wien, Bugra Suisse Wabern-Bern.
- Hanf, M. (1990) *The arable weeds of Europe*. BLV-Verlagsgesellschaft, Munich.
- Harper, J.L. (1977) *Population Biology of Plants*. Academic Press, New York.
- Harper, J.L. (1981) The population biology of modular organisms. In: *Theoretical ecology* (ed. R. May), pp. 53-77. Blackwell, Oxford.
- Haslam, S.M. (1972) *Phragmites communis* Trin. *Journal of Ecology*, 60, 585-610.
- Hawkes, J.C., Pyatt, D.G. & White, I.M.S. (1997) Using Ellenberg indicator values to assess soil quality in British forests from ground vegetation: a pilot study. *Journal of Applied Ecology*, 34, 375-387.
- Hegi, G. (1998). *Illustrierte Flora von Mitteleuropa*; I 3. Verlag Paul Parey, Berlin-Hamburg.
- Herrera, C.M., Jordano, P., Guitián, J. & Traveset, A. (1998) Annual variability in seed production by woody plants and the masting concept: reassessment of principles and relationship to pollination and seed dispersal. *American Naturalist*, 152, 576-594.
- Heyn, C.C., Dagan, O. & Nachman, B. (1974) The annual *Calendula* species, taxonomy and relationships. *Israel Journal of Botany*, 23, 169-201.
- Hill, M.O. & Carey, P.D. (1997) Prediction of yield in the Rothamstead Park grass experiment by Ellenberg indicator values. *Journal of Vegetation Science*, 8, 579-586.
- Hodgson, J.G. & Grime, J.P. (1990) The role of dispersal mechanisms, regeneration strategies and seed banks in the vegetation dynamics of the British landscape. In: *Species dispersal in Agricultural Habitats* (eds R.G.H. Bunce & D.C. Howard). Belhaven Press, London.
- Hodgson, J.G. (1986) Commonness and rarity in plants with special reference to the Sheffield Flora. Part II. The relative importance of climate, soils and land use. *Biological Conservation*, 36, 253-274.
- Hodgson, J.G., Grime, J.P., Hunt, R. & Thompson, K. (1995) *The Electronic Comparative Plant Ecology*. Chapman and Hall, London.
- Hodkinson, D.J., Askew, A.P., Thompson, K., Hodgson, J.G., Bakker, J.P. & Bekker, R.M. (1998) Ecological correlates of seed size in the British flora. *Functional Ecology*, 12, 762-766.
- Holm, T. (1925) On the development of buds upon roots and leaves. *Annals of Botany* 39, 867-881.
- Horvath, D.P. & Anderson, J.V. (2002) A molecular approach to understanding root bud dormancy in leafy spurge. *Weed Science*, 50, 227-231.
- Horvath, D.P., Chao, W.S. & Anderson, J.V. (2002) Molecular analysis of signals controlling dormancy and growth in underground adventitious buds of leafy spurge. *Plant Physiology*, 128, 1439-1446.
- Hughes, L., Dunlop, M., French, K., Leishman, M.R., Rice, B., Rodgerson, L. & Westoby, M. (1994) Predicting dispersal spectra - a minimal set of hypotheses based on plant attributes. *Journal of Ecology*, 82, 933-950.
- Huhta, A-P., Hellström, K., Rautio, P. & Tuomi, J. (2003) Grazing tolerance of *Gentianella amarella* and other monocarpic herbs: why is tolerance highest at low damage levels? *Plant Ecology*, 166, 49-61.
- Irmisch, T. (1850) *Zur Morphologie der monokotylyischen Knollen- und Zwiebelgewächse*. G. Reimer Verlag, Berlin.
- Iwasa, Y. & Kubo, T. (1997) Optimal size of storage for recovery after unpredictable disturbances. *Evolutionary Ecology*, 11, 41-65.
- Jackel, A.-K. & Poschlod, P. (1996) Plant strategies on continental dry grassland sites. *Verhandlungen Gesellschaft Ökologie*, 26, 511-517.
- Jackson, R.B. & Caldwell, M.M. (1993) The scale of nutrient heterogeneity around individual plants and its quantification with geostatistics. *Ecology*, 74, 612-614.
- Jäger, E.J. & K. Werner (2000): *Rothmaler Exkursionsflora von Deutschland. Gefäßpflanzen: Atlasband*. Fischer, Jena.
- Jäger, E.J. & Werner, K. (eds.) (2002) *Exkursionsflora von Deutschland. Band 4. Gefäßpflanzen: Kritischer Band*. Spektrum Akademischer Verlag, Heidelberg.

- Jakobsson, A. & Eriksson, O. (2000) A comparative study of seed number, seed size, seedling size and recruitment in grassland plants. *Oikos*, 88, 494-502.
- Janzen, D.H. (1984) Dispersal of small seeds by big herbivores: foliage is the fruit. *American Naturalist*, 123, 338-353.
- Jensen, H.A. (1998) *Bibliography on seed morphology*. A.A. Balkema, Rotterdam.
- Jones, E.W. (1959) Biological Flora of the British Isles: *Quercus* (L.). *Journal of Ecology*, 47, 169-222.
- Jongejans, E. & Schippers, P. (1999) Modelling seed dispersal by wind in herbaceous species. *Oikos*, 87, 362-272.
- Jónsdóttir, I.S. & Watson, M.A. (1997) Extensive physiological integration: An adaptive trait in resource-poor environments? *The ecology and evolution of clonal plants* (eds. H. de Kroon & J.M. van Groenendael), pp. 109-136. Backhuys Publishers, Leiden.
- Karrenberg, S., Edwards, P.J. & Kollmann, J. (2002). The life history of Salicaceae living in the active zone of floodplains. *Freshwater Biology*, 47, 733-748.
- Kelly, D. & Sork, V.L. (2002) Mast seeding in perennial plants: why, how, where? *Annual Review of Ecology and Systematics*, 33, 427-447.
- Kice (2002) What is terminal velocity? URL: <http://www.kice.com/multi-aspirator-background.htm>.
- Kidson, R. & Westoby, M. (2000) Seed mass and seedling dimensions in relation to seedling establishment. *Oecologia*, 125, 11-17.
- Kiviniemi, K. & O. Eriksson (1999) Dispersal, recruitment and site occupancy of grassland plants in fragmented habitats. *Oikos*, 86, 241-253.
- Kiviniemi, K. (1996) A study of adhesive seed dispersal of three species under natural conditions. *Acta Botanica Neerlandica*, 45, 73-83.
- Kjellsson, G. (1985) Seed fall and phenological overlap in a guild of ant-dispersed herbs. *Oecologia*, 68, 140-146.
- Klapp, E. (1965) *Grünlandvegetation und Standort*. Parey, Berlin.
- Kleyer, M. (1995) *Biological traits of vascular plants - a database*. University of Stuttgart, Arbeitsberichte, N.F. 2, Stuttgart.
- Klimeš, L. & Klimešová, J. (1999a) CLO-PLA2 - A database of clonal plants in central Europe. *Plant Ecology*, 141, 9-19.
- Klimeš, L. & Klimešová, J. (1999b) Root sprouting in *Rumex acetosella* under different nutrient levels. *Plant Ecology*, 141, 33-39.
- Klimeš, L. & Klimešová, J. (2000) Plant rarity and the mode of clonal growth. *Zeitschrift für Ökologie und Naturschutz*, 9, 43-52.
- Klimeš, L. & Klimešová, J. (2002) The effects of mowing and fertilization on carbohydrate reserves and regrowth of grasses: do they promote plant coexistence in species-rich meadows? *Evolutionary Ecology*, 15, 363-382.
- Klimeš, L., Klimešová, J., Hendriks, R. & Van Groenendael, J.M. (1997) Clonal plant architectures: a comparative analysis of form and function. *The ecology and evolution of clonal plants*. (eds. H. de Kroon & J.M. van Groenendael), pp. 1-29. Backhuys Publishers, Leiden.
- Klimeš, L. & Klimešová, J. (2003) Resprouting of herbs in disturbed habitats: is it adequately described by Bellingham-Sparrow's model? *Oikos*, 103, 225-229.
- Klimeš, L. & Klimešová, J. (XXXX) Root sprouting plants in central Europe. In prep.
- Klimešová, J. & Martínková, J. (2005) Intermediate growth forms as a model for study of plant clonality functioning: an example with root sprouters. *Evolutionary Ecology*, in press.
- Klimešová, J. (2001) Adventivní odnožování - přehlížená vlastnost kořenů rostlin. *Zprávy České Botanické Společnosti, Praha, Materiály*, 18, 63-72.
- Klimešová, J. (2003) Monokarpické rostliny schopné přežít silnou disturbanci (Monocarpic plants surviving severe disturbance). *Zprávy České Botanické Společnosti, Praha, Materiály*, in press.
- Klotz, S., Kühn, I. & Durka, W. (2002) BIOFLOR - Eine Datenbank mit biologisch-ökologischen Merkmalen zur Flora von Deutschland. *Schriftenreihe für Vegetationskunde*, 38, 1-334.
- Knevel, I.C., Bekker, R.M., Bakker, J.P. & Kleyer, M. (2003) Life-history traits of the Northwest European flora: The LEDA database. *Journal of Vegetation Science*, 14, 611-614.
- Koenig, W.D. & Knops, J.M.H. (2000) Patterns of annual seed production by northern hemisphere trees: a global perspective. *American Naturalist*, 155, 59-69.
- Koop, H. (1987) Vegetative reproduction of trees in some European natural forests. *Vegetatio*, 72, 103-110.
- Korpelainen, H. (1995) Mating system and distribution of enzyme genetic-variation in Bracken (*Pteridium aquilinum*). *Canadian Journal of Botany*, 73, 1611-1617.
- Korsmo, E. (1930) *Unkräuter im Ackerbau der Neuzeit*. Verlag von Julius Springer, Berlin.
- Kranz, O. (2000) Böden und Landnutzung auf Mauritius - Aspekte einer geoökologischen Detailuntersuchung mit Hilfe Geographischer Informationssysteme. MScDiplomarbeit am Fachbereich Geographie, Marburg.

- Kruijij, A.A., De Vries, D.M. & Mooi, H. (1967) Bijdrage tot de oecologie van Nederlandse graslandplanten. *Verslagen Landbouwkundige Onderzoeken* 696. Pudoc, Wageningen.
- Krumbiegel, A. (2001) Vegetative reproduction strategies of pseudo-annual plants in central Europe. *Beiträge zur Biologie der Pflanzen*, 72, 287-313.
- Kunzmann, D. (1993) *Vegetationskundliche und populationsbiologische Untersuchungen zum Blaustern (Scilla bifolia L.) am Oberrhein*. MSc Thesis University of Freiburg, Freiburg.
- Kunzmann, D. (2000) *Untersuchungen zur Diasporenbank und des Samenflugs trocken-magerer Standorte als Beitrag zur Definition biotischer Entwicklungspotentiale*. Dissertationes Botanicae Band 326, Gebrüder Borntraeger, Berlin, Stuttgart.
- Küppers, M. (1989) Hecken und Flurgehölze: Paradeobjekte für produktions- und populationsbiologische Untersuchungen an Holzgewachsen. *Verhandlungen der Gesellschaft für Ökologie*, 18, 689-700.
- Kutschera, L. & Lichtenegger, E. (1982) *Wurzelatlas mitteleuropäischer Grünlandpflanzen. Bd. 1, Monocotyledoneae*. Gustav Fischerverlag, Stuttgart.
- Kutschera, L. & Lichtenegger, E. (1992) *Wurzelatlas mitteleuropäischer Grünlandpflanzen. Teil 1, Band 2. Pteridophyta und Dicotyledonae (Magnoliopsida)*. Stuttgart, Fischer.
- Kutschera, L. & Lichtenegger, E. (1992) *Wurzelatlas mitteleuropäischer Grünlandpflanzen. Band 2: Pteridophyta und Dicotyledonae (Magnoliopsida). Teil 1: Morphologie, Anatomie, Ökologie, Verbreitung, Soziologie, Wirtschaft*. Gustav Fischer, Stuttgart.
- Landolt, E. (1977) Ökologische Zeigerwerte zur Schweizer Flora. *Veröffentlichungen des Geobotanischen Institutes der Eidg. Techn. Hochschule, Stiftung Rübél, Zürich* 64, 5-208.
- Lehsen, V. (2002) Lafore – Leaf area for everyone. URL: <http://www.uni-oldenburg.de/landeco/Download/Software/Laforem/Lafore.html>.
- Leishman, M.R. & Westoby, M. (1994a) The role of seed size in seedling establishment in dry soil conditions – experimental evidence from semi-arid species. *Journal of Ecology*, 82, 249-258.
- Leishman, M.R. & Westoby, M. (1994b) The role of large seed size in shaded conditions: experimental evidence. *Functional Ecology*, 8, 205-214.
- Leishman, M.R. (2001). Does the seed size/number trade-off model determine plant community structure? An assessment of the model mechanisms and their generality. *Oikos*, 93, 294-302.
- Lovett Doust, J. & Lovett Doust, L. (1988) *Plant Reproductive Ecology: Patterns and Strategies*. Oxford University Press, New York.
- Luftensteiner, H.W. (1982). *Untersuchungen zur Verbreitungsbiologie von Pflanzengemeinschaften an vier Standorten in Niederösterreich*. Bibliotheca Botanica, Vol. 135, Schweizerbartsche Buchhandlung, Stuttgart.
- Lukasiewicz, A. (1962) Morphologic development types of perennials. The Poznan Society of Friends of Science, Department of Mathematical and Natural Sciences. *Publication of the Section of Biology*, 27, 1-398. [in Polish]
- Martinková, J., Klimešová, J. & Mihulka, S. (2004a) Resprouting after disturbance: an experimental study with short-lived monocarpic herbs. *Folia Geobotanica*, 39, 1-12.
- Martinková, J., Kočvarová, M. & Klimešová, J. (2004b) Resprouting after disturbance in the short-lived herb *Rorippa palustris* (Brassicaceae): an experiment with juveniles. *Acta Oecologica*, 25, 143-150.
- McCartney, H.A. (1990) Dispersal mechanisms through the air. In: *Species Dispersal in Agricultural Habitats* (eds Bunce, R.G.H. & Howard, D.C.). pp. 133-158. Belhaven, London.
- Meyer, A.H. & Schmid, B. (1999) Experimental demography of the old-field perennial *Solidago altissima*: the dynamics of the shoot population. *Journal of Ecology*, 87, 17-27.
- Milton, W.E.J. (1948) The buried viable-seed content of upland soils in Montgomeryshire. *Empire Journal of Experimental Agriculture*, 16, 163-177.
- Morton, A. (2003) DMAP - Distribution mapping software. URL: <http://www.dmap.co.uk/>.
- Müller-Schneider, P. (1986) Verbreitungsbiologie der Blütenpflanzen Graubündens. *Veröffentlichungen des Geobotanischen Institutes der ETH, Stiftung Rübél*, 85, 263.
- Nathan, R., Katul, G.G., Horn, H.S., Thomas, S.M., Oren, R., Avissar, R., Pacala, S.W. & Levin, S.A. (2002) Mechanisms of long-distance dispersal of seeds by wind. *Nature*, 418, 409-413.
- Niklas, K.J. & Enquist, B.J. (2002) On the vegetative biomass partitioning of seed plant leaves, stems, and roots. *The American Naturalist*, 159, 482-497.
- Niklas, K.J. (1993) The allometry of plant reproductive biomass and stem diameter. *American Journal of Botany*, 80, 461-467.
- Niklas, K.J. (1994) *Plant allometry: the scaling of form and process*. The University of Chicago Press, Chicago.
- Niklas, K.J. (1995) Size-dependent Allometry of Tree Height, Diameter and Trunk-taper. *Annals of Botany*, 75, 217-227.

- Oborny, B., Kun, A., Czarán, T. & Bokros, S. (2000) The effect of clonal integration on plant competition for mosaic habitat space. *Ecology*, 81, 3291-3304.
- Otto, B. (2002) Merkmale von Samen, Früchten, generativen Germinulen und generativen Diasporen. In: BIOLFLORE – *Eine Datenbank mit biologisch-ökologischen Merkmalen zur Flora von Deutschland* (eds S. Klotz, I. Kühn, & W. Durka). Schriftenreihe für Vegetationskunde, 38, 177-196. Bundesamt für Naturschutz, Bonn-Bad Godesberg.
- Özer, Z. (1979) Über die Beeinflussung der Keimfähigkeit der Samen mancher Grünlandpflanzen beim Durchgang durch den Verdauungstrakt des Schafes und nach Mistgärung. *Weed Research*, 19, 247-254.
- Pärtel, M., Kalamees, R., Zobel, M. & Rosén, E. (1999) Alvar grasslands in Estonia: variation in species composition and community structure. *Journal of Vegetation Science*, 10, 561-570.
- Pärtel, M., Zobel, M., Zobel, K. & Van der Maarel, E. (1996) The species pool and its relation to species richness: evidence from Estonian plant communities. *Oikos*, 75, 111-117.
- Passioura, J.B. (1988) Water transport in and to roots. *Annual Review of Plant Physiology and Plant Molecular Biology*, 39, 245-265.
- Pavlu, V., Hejzman, M., Pavlu, L. & Gaisler, J. (2003) Effect of rotational and continuous grazing on vegetation of an upland grassland in the Jizerské Hory Mts., Czech Republic. *Folia Geobotanica*, 38, 21-34.
- Paz, H., Mazer, S.J. & Martinez-Ramos, M. (1999) Seed mass, seedling emergence, and environmental factors in seven rain forest Psychotria (Rubiaceae). *Ecology*, 80, 1594-1606.
- Peco, B., Traba, J., Levassor, C., Sanchez, A.M. & Azcarate, F.M. (2003) Seed size, shape and persistence in dry Mediterranean grass and scrublands. *Seed Science Research*, 13, 87-95.
- Peterson, R.L. (1975) The initiation and development of root buds. *The development and function of roots*. (eds J.G. Torrey & D.T. Clarkson), pp. 125-161. Academic Press, London.
- Piqueras, J. & Klimes, L. (1998) Clonal growth and population dynamics in the pseudo-annual plant *Trientalis europaea* L. *Plant Ecology*, 136, 213-227.
- Poschlod, P., Kleyer, M. & Tackenberg, O. (2000) Databases on life history traits as a tool for risk assessment in plant species. *Zeitschrift für Ökologie und Naturschutz*, 9, 3-18.
- Poschlod, P. & Jackel, A.-K. (1993) Untersuchungen zur Dynamik von generativen Diasporenbanken von Samenpflanzen in Kalkmagerrasen. I. Jahreszeitliche Dynamik des Diasporennetzes und der Diasporenbank auf zwei Kalkmagerrasenstandorten der Schwäbischen Alb. *Flora*, 188, 49-71.
- Pyankov, V. I., Kondratschuk, A.V. & Shipley, B. (1999) Leaf structure and specific leaf mass: the alpine desert plants of the Eastern Pamirs, Tadjikistan. *New Phytologist*, 143, 131-142.
- Rabotnov, T. A. (ed.) (1974) Biological flora of the Moscow region. Izdatelstvo Moskovskogo Universiteta, Moskva. [in Russian]
- Rauh, W. (1937) Die Bildung von Hypocotyl- und Wurzelsprossen und ihre Bedeutung für die Wuchsformen der Pflanzen. *Acta Nova Leopoldina, Halle*, 4/24, 395-555.
- Rauh, W. (1939) Über die Gesetzmäßigkeit der Verzweigung und deren Bedeutung für die Wuchsformen der Pflanzen. *Mitteilungen der deutschen dendrologischen Gesellschaft*, 52, 86-111.
- Raunkiaer, C. (1934) *The Life Forms of Plants and Statistical Plant Geography*. Clarendon, Oxford.
- Rejmánek, M. & Richardson, D. M. (1996) What attributes make some plant species more invasive? *Ecology*, 77, 1655-1661.
- Reyes, G., Brown, S., Chapman, J. and Lugo, A. E. (1992) *Wood densities of tropical tree species*. General Technical Report SO-88, US-Department of Agriculture, Forest Service, Southern Forest Experiment Station, New Orleans, USA.
- Richards, J.H. & Caldwell, M.M. (1985) Soluble carbohydrates, concurrent photosynthesis and efficiency in regrowth following defoliation: A field study with *Agropyron* species. *Journal of Applied Ecology*, 22, 907-920.
- Ridley, H. (1930) *The Dispersal of Plants throughout the World*. L. Reeve & Co. Ltd., Kent.
- Römermann, C., Tackenberg, O., Poschlod, P. (2005) How to predict attachment capacity potential of seeds to sheep and cattle coat from simple morphological traits. *Oikos*, 110, 219-230.
- Roo-Zielinska, E. & Solon, J. (1998) Phytosociological typology and bioindicator values of plant communities, as exemplified by meadows in the Nina valley, Southern Poland. *Documents Phytosociologiques*, 11, 543-554.
- Rothmaler, W. (2000) *Exkursionsflora von Deutschland*. Spektrum Akademischer Verlag, Heidelberg.
- Russell, E.W. (1973) *Soil Conditions and Plant Growth*. 10th Edition. Longman, London.
- Ryser P. (1996) The importance of tissue density for growth and life span of leaves and roots: a comparison of five ecologically contrasted grasses. *Functional Ecology*, 10, 717-723.
- Ryser, P. (1996) The importance of tissue density for growth and lifespan of leaves and roots: a comparison of five ecologically contrasting grasses. *Functional Ecology*, 10, 717-723.
- Salisbury, E.J. (1942) *The reproductive capacity of plants*. Bells, London.

- Sammul, M., Kull, K. & Tamm, A. (2003) Clonal growth in a species-rich grassland: Results of a 20-year fertilization experiment. *Folia Geobotanica*, 38, 1-20.
- Saverimuttu, T. & Westoby, M. (1996) Seedling longevity under deep shade in relation to seed size. *Journal of Ecology*, 84, 681-689.
- Schaffers, A.P. & Sýkora, K.V. (2000) Reliability of Ellenberg indicator values for moisture, nitrogen and soil reaction: a comparison with field measurements. *Journal of Vegetation Science* 11, 225-244.
- Schenk, H.J. (1999) Clonal splitting in desert shrubs. *Plant Ecology*, 141, 41-52.
- Sculthorpe, C.D. (1967) *The Biology of Aquatic Vascular Plants*. London.
- Šerá, B. & Šerý, M. (2004) Number and weight of seeds and reproductive strategies of herbaceous plants. *Folia Geobotanica*, 39, 27-40.
- Shipley, B. & Dion, J. (1992) The Allometry of seed production in herbaceous angiosperms. *The American Naturalist*, 139, 467-483.
- Shipley, B. & Vu, T.T. (2002) Dry matter content as a measure of dry matter concentration in plants and their parts. *New Phytologist*, 153, 359-364.
- Silvertown, J.W. & Lovett Doust, J. (1993) *Introduction to plant population biology*. Blackwell Scientific Publications, London.
- Stace, C.A. (1997) *New Flora of the British Isles*. 2nd edition. Cambridge University Press, Cambridge.
- Stöcklin, J. & Fischer, M. (1999) Plants with longer-lived seeds have lower local extinction rates in grassland remnants 1950-1985. *Oecologia*, 120, 539-543.
- Symonides, E. (1978) Numbers, distribution and specific composition of diaspores in the soils of the plant association Spergulo-Corynephoretum. *Ekologia Polska*, 26, 11-122.
- Tackenberg, O. (2003) Modelling long-distance dispersal of plant diaspores by wind. *Ecological Monographs*, 73, 173-189.
- Tackenberg, O., Poschlod, P. & Bonn, S. (2003) Assessment of wind dispersal potential in plant species. *Ecological Monographs*, 73, 191-205.
- Tallowin, J.R.B. & Smith, R.E.N. (2001) Restoration of a Cirsio-Molinietum fen meadow on an agriculturally improved pasture. *Restoration Ecology*, 9, 167-178.
- Talmon, E. (2002) Ausbreitung von Pflanzenarten-Modellierung der Ausbreitung von Pflanzenarten in Trockenstandorten. In: MOSAIK-Projekt - Halboffene Weide- und Felsgraslandschaften als alternative Leitbilder für die Pflege von Trockenstandorten und ihre Konsequenzen für das Überleben von Flora und Fauna. pp. 55-64. Teilprojekte MR2, University of Regensburg.
- Tamm, A., Kull, K. & Sammul, M. (2001) Classifying clonal growth forms based on vegetative mobility and ramet longevity: a whole community analysis. *Evolutionary Ecology*, 15, 383-401.
- Ter Braak, C.J.F. & Gremmen, N.J.M. (1987) Ecological amplitudes of plant species and the internal consistency of Ellenberg's indicator values for moisture. *Vegetatio*, 69, 79-87.
- Ter Heerdt, G.N.J., Verweij, G.L., Bekker, R.M. & Bakker, J.P. (1996) An improved method for seed bank analysis: seedling emergence after removing the soil by sieving. *Functional Ecology*, 10, 144-151.
- Thimonier, A., Dupouey, J.L., Bost, F. & Becker, M. (1994) Simultaneous eutrophication and acidification of a forest ecosystem in North-East France. *New Phytologist*, 126, 533-539.
- Thompson, K. & Grime, J.P. (1979) Seasonal variation in the seed banks of herbaceous species in ten contrasting habitats. *Journal of Ecology*, 67, 893-921.
- Thompson, K. (1992) The Functional Ecology of seed banks. In *Seeds: the Ecology of Regeneration in Plant Communities*. (ed. M. Fenner), pp. 231-258. CAB International, Wallingford.
- Thompson, K., Bakker, J.P. & Bekker, R.M. (1997) *The soil seed banks of North West Europe: methodology, density and longevity*. Cambridge University Press, Cambridge.
- Thompson, K., Bakker, J.P., Bekker, R.M. & Hodgson, J.G. (1998) Ecological correlates of seed persistence in soil in the north-west European flora. *Journal of Ecology*, 86, 163-169.
- Thompson, K., Band, S.R. & Hodgson, J.G. (1993) Seed size and shape predict persistence in soil. *Journal of Ecology*, 7, 236-241.
- Thompson, K., Jalili, A., Hodgson, J.G., Hamzeh'ee, B., Asri, Y., Shaw, S., Shirvany, A., Yazdani, S., Khoshnevis, M., Zarrinkamar, M.A. & Safavi, R. (2001) Seed size, shape and persistence in the soil in an Iranian flora. *Seed Science Research*, 11, 345-355.
- Tilman, D. (1988). *Plant strategies and the dynamics and structure of plant communities*. Princeton University Press, Princeton, NJ.
- Troll, W. (1935) *Vergleichende Morphologie der höheren Pflanzen: Vegetationsorgane*. Band 1, Teil 1, Berlin.
- Troll, W. (1937-1941) *Vergleichende Morphologie der höheren Pflanzen. Band 1 - Vegetationsorgane*. Verlag von Gebrüder Borntraeger, Berlin - Zahlendorf.
- USDA (Soil Service Staff) (1993). *Soil Survey Manual*. USDA - Soil Conservation Service, Agricultural Handbook No. 18, U.S. Gov. Print. Office, Washington. Download available at <http://soils.usda.gov/scientists.html>.



- Van Altena, S.C. & Minderhoud, J.W. (1972) Keimfähige Samen von Gräsern und Kräutern in der Narbeschicht der Niederländischen Weiden. *Zeitschrift für Acker- und Pflanzenbau*, 136, 95-109.
- Van der Maarel, E., Boot, R., Van Dorp, D. & Rijntjes, J. (1985) Vegetation succession on the dunes near Oostvoorne, The Netherlands; a comparison of the vegetation in 1959 and 1980. *Vegetatio*, 58, 137-187.
- Van der Maarel, E. (1993) Relations between sociological-ecological species groups and Ellenberg indicator values. *Phytocoenologia* 23, 277-289.
- Van der Pijl, L. (1982) Principles of dispersal in higher plants. Springer, Berlin.
- Verburg, R.G. (1998) The witch at the forest floor. Reproductive allocation and clonal diversity in the pseudo-annual *Circaea lutetiana* L. PhD thesis, Utrecht University, Utrecht.
- Vevle, O. & Aase, K. (1980) On the use of ecological factor numbers in Norwegian forest communities. *K. Nor. Vidensk. Selsk. Mus. Rapp. Bot. Ser.*, 5, 178-201.
- Vickery, J.A., Tallowin, J.R.B., Ferber, R.E., Asteraki, E.J., Atkinson, P.W., Fuller, R.J. & Brown, V.K. (2001) The management of lowland neutral grasslands in Britain: effects of agricultural practices on birds and their food resources. *Journal of Applied Ecology*, 38, 647-664.
- Wagenitz, G. (1996) *Wörterbuch der Botanik*. Gustav Fisher, Jena.
- Waller, D.M. & Steingraeber, D.A. (1995) Opportunities and Constraints in the Placement of Flowers and Fruits in the placement of Flowers and Fruits. In: *Plant stems: Physiology and Functional Morphology* (ed. B.L. Gartner), pp. 51-74. Academic Press, San Diego.
- Wehsarg, O. (1954) *Biologie der Ackerunkräuter*. Akademie - Verlag GmbH., Berlin.
- Weiher, E., Van der Werf, A., Thompson, K., Roderick, M., Garnier, E. & Eriksson, O. (1999) Challenging Theophrastus: A common core list of plant traits for *Functional Ecology*. *Journal of Vegetation Science*, 10, 609-620.
- Weiner, J. (1988) The Influence of Competition on Plant Reproduction. In: *Plant Reproductive Ecology: Patterns and Strategies* (eds. J. Lovett Doust & L. Lovett Doust.), pp. 228-245. Oxford University Press, New York.
- Westoby, M. (1998) A leaf-height-seed (LHS) plant ecology strategy scheme. *Plant and Soil*, 199, 213-227.
- Westoby, M., Leishman, M., Lord, J. (1997) Comparative ecology of seed size and dispersal. In: *Plant life histories* (Eds Silvertown, J., Franco, M., Harper, J.L.). pp. 143-162. Cambridge University Press, Cambridge.
- Wetzel, R.G. (2001) *Limnology, lake and river ecosystems*. Academic Press, New York.
- Wilson, B.F. (1995) Shrub stems: Form and function. In: *Plant stems: Physiology and Functional Morphology*. (ed. B.L. Gartner), pp. 91-102. Academic Press, San Diego.
- Wilson, P.J., Thompson, K. & Hodgson, J.G. (1999) SLA and leaf dry matter content as alternative predictors of plant strategies. *New Phytologist*, 143, 155-162.
- Wilson, P.J., Thompson, K. & Hodgson, J.G. (1999) Specific leaf area and leaf dry matter content as alternative predictors of plant strategies. *New Phytologist*, 143, 155-162.
- Wittrock, V.B. (1884a) Über Wurzelsprossen bei krautgarten Gewächsen, mit besonderer Rücksicht auf ihre verschiedene biologische Bedeutung. *Botanisches Centralblatt*, 17(8), 227-232.
- Wittrock, V.B. (1884b) Über Wurzelsprossen bei krautgarten Gewächsen, mit besonderer Rücksicht auf ihre verschiedene biologische Bedeutung. *Botanisches Centralblatt*, 17(9), 257-264.
- Wright, I.J. & Westoby, M. (1999) Differences in seedling growth behaviour among species: trait correlations across species, and trait shifts along nutrient compared to rainfall gradients. *Journal of Ecology*, 87, 85-97.
- Wright, I.J., Westoby, M. & Reich, P.B. (2002) Convergence towards higher leaf mass per area in dry and nutrient-poor habitats has different consequences for leaf lifespan. *Journal of Ecology*, 90, 534-543.
- Young, J.A. & Young, C.G. (1992) *Seeds of woody plants in North America*. Revised and enlarged edition. Dioscorides Press, Portland.
- Ziegenspeck, H. (1936) Orchidaceae. In: *Lebensgeschichte der Blütenpflanzen Mitteleuropas*. (eds. O. Kirschner, E. Loew & C. Schröter) pp. 1-840. Eugen Ulmer Verlag, Stuttgart.
- Zobler, L. (1986) *A world soil file for global climate modelling*. NASA TM-87802. National Aeronautics and Space Administration. Washington.
- Zophi, H-J. (1993a) Ecotypic variation in *Rhinanthus alectorolophus* (Scopoli) Pollich (Scrophulariaceae) in relation to grassland management I. Morphological delimitation's and habitats of seasonal ecotypes. *Flora*, 188, 15-39.
- Zophi, H-J. (1993b) Ecotypic variation in *Rhinanthus alectorolophus* (Scopoli) Pollich (Scrophulariaceae) in relation to grassland management II. The genotypic basis of seasonal ecotypes. *Flora*, 188, 153-173.
- Zophi, H-J. (1997) Ecotypic variation of *Euphrasia rostkoviana* Hayne (Scrophulariaceae) in relation to grassland management II. The genotypic basis of seasonal ecotypes. *Flora*, 192, 279-295.